

# Linear Models and Experimental Designs

STATS 752

Winter 2023

Cameron Roopnarine\*

Shui Feng<sup>†</sup>

9th April 2023

## Contents

|    |  |    |
|----|--|----|
| 1  | Lecture 1: Review of Linear Algebra                                  | 3  |
| 2  | Lecture 2: Quadratic Forms and Distributions                         | 8  |
| 3  | Lecture 3: Some Basic Lemmas   | 13 |
| 4  | Lecture 4: Quadratic Forms with Idempotency                          | 18 |
| 5  | Lecture 5: Criteria for Independence                                 | 22 |
| 6  | Lecture 6: Cochran's Theorem   | 25 |
| 7  | Lecture 7: Full Rank Regression                                      | 28 |
| 8  | Lecture 8: Test of Overall Regression                                | 32 |
| 9  | Lecture 9: Lack of Fit   | 37 |
| 10 | Lecture 10: Determination of Predictors and Generalized Inverse      | 39 |
| 11 | Lecture 11: $g$ -inverse   | 42 |
| 12 | Lecture 12: Regression Without Full Rank                             | 46 |
| 13 | Lecture 13: Regression Without Full Rank (Continued)                 | 49 |
| 14 | Lecture 14: General Linear Hypothesis Testing                        | 53 |
| 15 | Lecture 15: ANOVA Table For One-Way Classification                   | 58 |
| 16 | Lecture 16: Two-Way Nested Classification Part II                    | 62 |
| 17 | Lecture 17: Two-way Crossed Classification (No Interaction) Part I   | 65 |
| 18 | Lecture 18: Two-way Crossed Classification (No Interaction) Part II  | 68 |
| 19 | Lecture 19: Two-way Crossed Classification (No Interaction) Part III | 71 |

---

\* $\LaTeX$ er

<sup>†</sup>Instructor

|   |           |
|---|-----------|
| <b>20 Lecture 20: Two-way Crossed Classification (No Interaction) Part IV</b> | <b>74</b> |
| <b>21 Lecture 21: Two-way Crossed Classification (Interaction) Part II</b>    | <b>77</b> |
| <b>22 Lecture 22: Factorial Design</b>  | <b>80</b> |

# 1 Lecture 1: Review of Linear Algebra

## DEFINITION 1.1: Vectors in $\mathbb{R}^n$

For any positive integer  $n$ , we define  $n$ -dimensional Euclidean space  $\mathbb{R}^n$  by

$$\mathbb{R}^n = \left\{ \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \mid x_1, \dots, x_n \in \mathbb{R} \right\}.$$

If  $\vec{x} \in \mathbb{R}^n$ , then there exists  $x_1, \dots, x_n \in \mathbb{R}$  such that

$$\vec{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}.$$

## DEFINITION 1.2: Matrix

An  $n \times m$  **matrix**  $\mathbf{A}$  is a rectangular array with  $n$  rows and  $m$  columns. We denote the entry in the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column by  $a_{ij}$  or  $(\mathbf{A})_{ij}$ . That is,

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1j} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2j} & \cdots & a_{2m} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{im} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nj} & \cdots & a_{nm} \end{pmatrix}.$$

The set of all  $n \times m$  matrices with real entries is denoted by  $\mathbb{R}^{n \times m}$ .

## DEFINITION 1.3: Transpose

Let  $\mathbf{A} \in \mathbb{R}^{n \times m}$ . We define the **transpose** of  $\mathbf{A}$ , denoted  $\mathbf{A}'$ , to be the  $m \times n$  matrix whose  $ij^{\text{th}}$  entry is the  $ji^{\text{th}}$  entry of  $\mathbf{A}$ . That is,

$$(\mathbf{A}')_{ij} = (\mathbf{A})_{ji}.$$

## DEFINITION 1.4: Square Matrix

An  $n \times n$  matrix is called a **square matrix**.

## DEFINITION 1.5: Symmetric Matrix

A matrix is called **symmetric** if  $\mathbf{A}' = \mathbf{A}$ .

## DEFINITION 1.6: Diagonal Matrix

An  $n \times n$  matrix  $\mathbf{D}$  is said to be a **diagonal matrix** if  $d_{ij} = 0$  for all  $i \neq j$ . We denote a diagonal matrix by

$$\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n).$$

**DEFINITION 1.7: Identity Matrix**

The  $n \times n$  matrix  $\mathbf{I}$  (or  $\mathbf{I}_n$ ) such that  $(\mathbf{I})_{ii} = 1$  for  $1 \leq i \leq n$ , and  $(\mathbf{I})_{ij} = 0$  whenever  $i \neq j$  is called the **identity matrix**.

**DEFINITION 1.8: Upper Triangular, Lower Triangular**

An  $n \times m$  matrix  $\mathbf{U}$  is said to be **upper triangular** if  $u_{ij} = 0$  whenever  $i > j$ . An  $n \times m$  matrix  $\mathbf{L}$  is said to be **lower triangular** if  $\ell_{ij} = 0$  whenever  $i < j$ .

- Upper triangular:

$$\mathbf{U} = \begin{pmatrix} u_{11} & u_{12} & \cdots & u_{1m} \\ 0 & u_{22} & \ddots & \vdots \\ \vdots & \ddots & \ddots & u_{(n-1)m} \\ 0 & \cdots & 0 & u_{nm} \end{pmatrix}.$$

- Lower triangular:

$$\mathbf{L} = \begin{pmatrix} \ell_{11} & 0 & \cdots & 0 \\ \ell_{21} & \ell_{22} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ \ell_{n1} & \cdots & \ell_{n(m-1)} & \ell_{nm} \end{pmatrix}.$$

**DEFINITION 1.9: Vector/Matrix of 1's and 0's**

Vector of 1's:

$$\vec{j} = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^n.$$

Matrix of 1's:

$$\mathbf{J} = \begin{pmatrix} 1 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 1 \end{pmatrix} \in \mathbb{R}^{n \times n}.$$

Zero vector:

$$\vec{0} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix} \in \mathbb{R}^n.$$

Zero Matrix:

$$\mathbf{O} = \begin{pmatrix} 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{pmatrix} \in \mathbb{R}^{n \times n}.$$

**DEFINITION 1.10: Matrix Mapping**

If  $\mathbf{A}$  is an  $n \times m$  matrix, then we can define a function  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  by  $T(\vec{x}) = \mathbf{A}\vec{x}$  called a **matrix mapping**. For this mapping, we define:

- Kernel

$$\text{Ker}(T) = \{\vec{x} \in \mathbb{R}^m \mid \mathbf{A}\vec{x} = \vec{0}\}.$$

- **Image**

$$\text{Image}(T) = \{\mathbf{A}\vec{x} \in \mathbb{R}^n \mid \vec{x} \in \mathbb{R}^m\}.$$

- **Rank**

$$\text{rank}(T) = \dim(\text{Image}(T)).$$

- **Nullity**

$$\text{nullity}(T) = \dim(\text{Ker}(T)).$$

**REMARK — Rank-Nullity Theorem**

$$\text{rank}(T) + \text{nullity}(T) = m.$$

**REMARK**

Note that

$$\mathbf{A} = \begin{pmatrix} a_{11} & \cdots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nm} \end{pmatrix} = \begin{pmatrix} \vec{a}_1 \\ \vdots \\ \vec{a}_n \end{pmatrix} = (\vec{a}^1 \ \cdots \ \vec{a}^m).$$

Clearly, the image of  $T$  is the space generated by  $\vec{a}^1, \dots, \vec{a}^m$ . Therefore,

$$\text{rank}(T) = \text{column rank of } \mathbf{A}.$$

**THEOREM 1.1**

$\mathbf{A}$  and  $\mathbf{A}'$  have the same column rank.

**Proof:** Let  $\mathbf{A} \rightarrow T$ ,  $\mathbf{A}'\mathbf{A} \rightarrow \tilde{T}$ , and  $\mathbf{A}' \rightarrow \hat{T}$ .

(1) Let  $\vec{x} \in \text{Ker}(T)$ , so we have  $\mathbf{A}\vec{x} = \vec{0} \implies \mathbf{A}'\mathbf{A}\vec{x} = \vec{0}$ . Hence,  $\vec{x} \in \text{Ker}(\tilde{T})$ . So,  $\text{Ker}(T) \subset \text{Ker}(\tilde{T})$ .

(2)  $\mathbf{A}'\mathbf{A}\vec{x} = \vec{0} \implies \vec{x}'\mathbf{A}'\mathbf{A}\vec{x} = 0 \implies (\mathbf{A}\vec{x})'\mathbf{A}\vec{x} = 0 \implies \mathbf{A}\vec{x} = \vec{0}$ . So,  $\text{Ker}(\tilde{T}) \subset \text{Ker}(T)$ .

Therefore,  $\text{rank}(T) = \text{rank}(\tilde{T})$ . By Rank-Nullity theorem,  $\mathbf{A}$  and  $\mathbf{A}'$  have the same column rank, noting that  $\text{Image}(\tilde{T}) \subset \text{Image}(\hat{T})$ . Hence, the column rank of  $\mathbf{A}$  is less than or equal to the column rank of  $\mathbf{A}'$ . By symmetry, the column rank of  $\mathbf{A}'$  is less than or equal to the column rank of  $(\mathbf{A}')' = \mathbf{A}$ . Therefore,  $\mathbf{A}$  and  $\mathbf{A}'$  have the same column rank.

**REMARK**

The column rank of  $\mathbf{A}'$  is the row rank of  $\mathbf{A}$ . Hence,

$$\begin{aligned} \text{rank}(\mathbf{A}) &= \text{maximum number of linearly independent rows of } \mathbf{A} \\ &= \text{maximum number of linearly independent columns of } \mathbf{A}. \end{aligned}$$

**DEFINITION 1.11: Full Rank**

Let  $\mathbf{A} \in \mathbb{R}^{n \times m}$ . We say  $\mathbf{A}$  has **full rank** if  $\text{rank}(\mathbf{A}) = \min\{n, m\}$

**THEOREM 1.2**

Let  $\mathbf{A} \in \mathbb{R}^{n \times m}$  and  $\mathbf{B} \in \mathbb{R}^{m \times p}$ .

- (1)  $(\mathbf{AB})' = \mathbf{B}'\mathbf{A}' \in \mathbb{R}^{p \times n}$ .
- (2)  $\vec{j}'\vec{j} = n$  and  $\vec{j}\vec{j}' = \mathbf{J}$ .
- (3)  $\mathbf{J}\mathbf{J} = n\mathbf{J}$ .

**THEOREM 1.3**

- (1)  $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}')$ .
- (2)  $\text{rank}(\mathbf{A}'\mathbf{A}) = \text{rank}(\mathbf{A}\mathbf{A}') = \text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}')$ .
- (3)  $\text{rank}(\mathbf{AB}) \leq \min\{\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B})\}$ .

**Proof:** We have already shown (1) and (2). For (3), we have

$$\text{rank}(\mathbf{AB}) \leq \text{rank}(\mathbf{A}).$$

On the other hand,

$$\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{B}'\mathbf{A}') \leq \text{rank}(\mathbf{B}') = \text{rank}(\mathbf{B}).$$

**REMARK — Invertible Matrix Theorem**

$\mathbf{A} \in \mathbb{R}^{n \times n}$  is invertible (non-singular) if and only if  $\text{rank}(\mathbf{A}) = n$ , and we denote the inverse of  $\mathbf{A}$  by  $\mathbf{A}^{-1}$ .

**REMARK — Properties of Invertible Matrices**

- (1)  $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$ .
- (2)  $(\mathbf{A}')^{-1} = (\mathbf{A}^{-1})'$ .
- (3)  $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$ .
- (4)  $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$ .

**DEFINITION 1.12: Positive Definite, Positive Semidefinite**

- $\mathbf{A}$  is **positive definite** when  $\vec{x}'\mathbf{A}\vec{x} > 0 \iff \vec{x} \neq \vec{0}$ .
- $\mathbf{A}$  is **positive semidefinite** when  $\vec{x}'\mathbf{A}\vec{x} \geq 0$  for all  $\vec{x}$  and there exists  $\vec{x} \neq \vec{0}$  such that  $\vec{x}'\mathbf{A}\vec{x} = 0$ .

**DEFINITION 1.13: Orthogonal**

$\mathbf{A} \in \mathbb{R}^{n \times n}$  is **orthogonal** if  $\mathbf{A}' = \mathbf{A}^{-1}$ .

**DEFINITION 1.14: Eigenvalue, Eigenvector, Spectrum**

Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$ . If there exists a vector  $\vec{x} \neq \vec{0}$  such that  $\mathbf{A}\vec{x} = \lambda\vec{x}$ , then  $\lambda$  is called an **eigenvalue** of  $\mathbf{A}$  and  $\vec{x}$  is called an **eigenvector** of  $\mathbf{A}$  corresponding to  $\lambda$ . The set of all eigenvalues of  $\mathbf{A}$  is called the **spectrum** for  $\mathbf{A}$ .

**EXAMPLE 1.1**

$$\underbrace{\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}}_{\mathbf{A}} \underbrace{\begin{pmatrix} 1 \\ 1 \end{pmatrix}}_{\vec{v}} = \underbrace{3}_{\lambda} \underbrace{\begin{pmatrix} 1 \\ 1 \end{pmatrix}}_{\vec{v}}.$$

**THEOREM 1.4: Spectral Decomposition**

Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$ .  $\mathbf{A}$  is symmetric if and only if

$$\mathbf{A} = \mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n) \mathbf{Q},$$

where  $\mathbf{Q}$  is an orthogonal matrix, that is,  $\mathbf{Q}\mathbf{Q}' = \mathbf{I}$ .

**THEOREM 1.5**

Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$  be a symmetric matrix.

- (i)  $\mathbf{A}$  is positive definite (semidefinite) if and only if all eigenvalues are positive (non-negative).
- (ii)  $\mathbf{A}$  is positive definite if and only if there exists a unique lower triangular matrix  $\mathbf{L}$  with positive diagonal elements such that  $\mathbf{A} = \mathbf{L}\mathbf{L}'$  (Cholesky decomposition).

**DEFINITION 1.15: Idempotent, Trace**

- $\mathbf{A} \in \mathbb{R}^{n \times n}$  is **idempotent** if  $\mathbf{A} = \mathbf{A}^2$ .
- Let  $\text{tr}: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$  be defined by

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^n a_{ii}$$

(called the **trace** of a matrix).

**THEOREM 1.6**

Let  $\mathbf{A} \in \mathbb{R}^{n \times m}$ ,  $\mathbf{B} \in \mathbb{R}^{m \times p}$ ,  $\mathbf{C} \in \mathbb{R}^{p \times n}$ , and  $a, b \in \mathbb{R}$ .

- (1)  $\text{tr}$  is linear:  $\text{tr}(a\mathbf{A} + b\mathbf{B}) = a \text{tr}(\mathbf{A}) + b \text{tr}(\mathbf{B})$ .
- (2) Cyclic property:  $\text{tr}(\mathbf{ABC}) = \text{tr}(\mathbf{CAB}) = \text{tr}(\mathbf{BCA})$ .

**Proof:**

- (1) By definition.

(2) Note that

$$\begin{aligned}
 \text{tr}(\mathbf{ABC}) &= \sum_{i=1}^n \sum_{k=1}^m \sum_{j=1}^p a_{ik} b_{kj} c_{ji} \\
 &= \sum_{j=1}^p \sum_{i=1}^n \sum_{k=1}^m c_{ji} a_{ik} b_{kj} \\
 &= \text{tr}(\mathbf{CAB}) \\
 &= \sum_{k=1}^m \sum_{j=1}^p \sum_{i=1}^n b_{kj} c_{ji} a_{ik} \\
 &= \text{tr}(\mathbf{BCA}).
 \end{aligned}$$

### REMARK — Properties of Idempotent Matrices

(1) Eigenvalues of idempotent matrices are 1 or 0 since

$$\begin{aligned}
 \mathbf{A}\vec{x} &= \lambda\vec{x} \\
 \implies \mathbf{AA}\vec{x} &= \lambda\mathbf{A}\vec{x} = \lambda^2\vec{x} \\
 \implies \lambda\vec{x} &= \lambda^2\vec{x} \\
 \implies \lambda &= \lambda^2 \\
 \implies \lambda &= 0 \text{ or } 1.
 \end{aligned}$$

(2) Idempotent matrices are diagonalizable, that is, there exists an invertible matrix  $\mathbf{P}$  such that  $\mathbf{A} = \mathbf{P}^{-1}\mathbf{D}\mathbf{P}$ , where

$$\mathbf{D} = \text{diag}(\lambda_1, \dots, \lambda_n), \quad \forall \lambda_i = 0 \text{ or } 1.$$

(3) If  $\mathbf{A}$  is idempotent, then  $\text{tr}(\mathbf{A}) = \text{rank}(\mathbf{A})$ .

$$\begin{aligned}
 \text{tr}(\mathbf{A}) &= \text{tr}(\mathbf{P}^{-1}\mathbf{D}\mathbf{P}) \\
 &= \text{tr}(\mathbf{D}\mathbf{P}\mathbf{P}^{-1}) \\
 &= \text{tr}(\mathbf{D}) \\
 &= \lambda_1 + \dots + \lambda_n \\
 &= \# \text{ of non-zero } \lambda_i\text{'s} \\
 &= \text{rank}(\mathbf{A}).
 \end{aligned}$$

LECTURE 2  
12th January

## 2 Lecture 2: Quadratic Forms and Distributions

### DEFINITION 2.1: Quadratic Form

The **quadratic form** associated  $\mathbf{A} \in \mathbb{R}^{n \times n}$  is defined as

$$\vec{x}'\mathbf{A}\vec{x} = \sum_{i=1}^n \sum_{j=1}^n x_i a_{ij} x_j.$$



If  $\tilde{\mathbf{A}} = \frac{\mathbf{A} + \mathbf{A}'}{2}$ , then note that  $\tilde{\mathbf{A}}$  is symmetric and

$$\begin{aligned}\vec{x}' \tilde{\mathbf{A}} \vec{x} &= \sum_{i=1}^n \sum_{j=1}^n x_j \frac{a_{ij} + a_{ji}}{2} x_i \\ &= \sum_{i=1}^n \sum_{j=1}^n x_i a_{ij} x_j \\ &= \vec{x}' \mathbf{A} \vec{x}.\end{aligned}$$

Therefore, there is a one-to-one correspondence between quadratic forms and symmetric matrices.

### EXAMPLE 2.1

Let  $S^2$  denote the sample variance of a random sample  $X_1, \dots, X_n$ . Set

$$\vec{X} = \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}, \quad \vec{\mu} = \mathbb{E}[\vec{X}] = \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_n \end{pmatrix}.$$

Write  $(n-1)S^2$  as a quadratic form and identify the matrix  $\mathbf{A}$ .

**Solution:**

$$\begin{aligned}(n-1)S^2 &= \sum_{i=1}^n (X_i - \bar{X})^2 \\ &= \sum_{i=1}^n X_i^2 - 2 \left( \sum_{i=1}^n X_i \right) \bar{X} + n\bar{X}^2 \\ &= \sum_{i=1}^n X_i^2 - \frac{1}{n} \left( \sum_{i=1}^n X_i \right)^2 \\ &= \vec{X}' \vec{X} - \frac{1}{n} (\vec{j}' \vec{X})^2 \\ &= \vec{X}' \vec{X} - \frac{1}{n} \vec{X}' \vec{j} \vec{j}' \vec{X} \\ &= \vec{X}' \left( \mathbf{I} - \frac{1}{n} \vec{j} \vec{j}' \right) \vec{X} \\ &= \vec{X}' \left( \mathbf{I} - \frac{1}{n} \mathbf{J} \right) \vec{X}.\end{aligned}$$

Hence,  $\mathbf{A} = \mathbf{I} - \frac{1}{n} \mathbf{J}$  is symmetric.

### THEOREM 2.1

If  $\vec{X}$  is a random vector with mean  $\vec{\mu}$ , covariance matrix  $\Sigma$ , and  $\mathbf{A}$  is a symmetric matrix of constants, then

$$\mathbb{E}[\vec{X}' \mathbf{A} \vec{X}] = \text{tr}(\mathbf{A} \Sigma) + \vec{\mu}' \mathbf{A} \vec{\mu}.$$

**Proof:**

$$\begin{aligned}
\mathbb{E}[\vec{X}'\mathbf{A}\vec{X}] &= \mathbb{E}[\text{tr}(\vec{X}'\mathbf{A}\vec{X})] \\
&= \mathbb{E}[\text{tr}(\mathbf{A}\vec{X}\vec{X}')] \\
&= \text{tr}(\mathbb{E}[\mathbf{A}\vec{X}\vec{X}']) \\
&= \text{tr}(\mathbf{A}\mathbb{E}[\vec{X}\vec{X}']) \\
&= \text{tr}\left(\mathbf{A}\mathbb{E}[(\vec{X} - \vec{\mu} + \vec{\mu})(\vec{X}' - \vec{\mu}' + \vec{\mu}')]\right) \\
&= \text{tr}(\mathbf{A}[\boldsymbol{\Sigma} + \vec{\mu}\vec{\mu}']) \\
&= \text{tr}(\mathbf{A}\boldsymbol{\Sigma}) + \text{tr}(\mathbf{A}\vec{\mu}\vec{\mu}') \\
&= \text{tr}(\mathbf{A}\boldsymbol{\Sigma}) + \text{tr}(\vec{\mu}'\mathbf{A}\vec{\mu}) \\
&= \text{tr}(\mathbf{A}\boldsymbol{\Sigma}) + \vec{\mu}'\mathbf{A}\vec{\mu}.
\end{aligned}$$

**EXAMPLE 2.2**

Assume that  $\mu_1 = \dots = \mu_n = \mu$  and  $\boldsymbol{\Sigma} = \sigma^2\mathbf{I}$ . Find  $\mathbb{E}[S^2]$ .

**Solution:**

$$\begin{aligned}
\mathbb{E}[S^2] &= \frac{1}{n-1} \mathbb{E}[(n-1)S^2] \\
&= \frac{1}{n-1} \mathbb{E}\left[\vec{X}'(\mathbf{I} - \frac{1}{n}\mathbf{J})\vec{X}\right] \\
&= \frac{1}{n-1} \mathbb{E}\left[\text{tr}\left((\mathbf{I} - \frac{1}{n}\mathbf{J})\boldsymbol{\Sigma}\right) + \vec{\mu}'(\mathbf{I} - \frac{1}{n}\mathbf{J})\vec{\mu}\right] \\
&= \frac{1}{n-1} \mathbb{E}\left[\text{tr}\left((\mathbf{I} - \frac{1}{n}\mathbf{J})\sigma^2\mathbf{I}\right) + \mu^2\vec{j}'(\mathbf{I} - \frac{1}{n}\mathbf{J})\vec{j}\right] \\
&= \frac{1}{n-1} \mathbb{E}\left[\sigma^2(n-1) + \mu^2(\vec{j}'\vec{j} - \frac{1}{n}\vec{j}'\mathbf{J}\vec{j})\right] \\
&= \frac{1}{n-1} \mathbb{E}[\sigma^2(n-1) + 0] \qquad \text{since } \vec{j}'\mathbf{J}\vec{j} = n\vec{j}'\vec{j} \\
&= \sigma^2.
\end{aligned}$$

**REMARK — Multivariate Normal Distribution**

Let  $\vec{X} = (X_1, X_2, \dots, X_n)$  be a  $1 \times n$  random vector with  $\mathbb{E}[X_i] = \mu_i$  and  $\text{Cov}(X_i, X_j) = \sigma_{ij}$ , for  $i, j = 1, 2, \dots, n$ . Let  $\vec{\mu} = (\mu_1, \mu_2, \dots, \mu_n)$  be the mean vector and  $\boldsymbol{\Sigma}$  be the  $n \times n$  symmetric covariance matrix whose  $(i, j)$  entry is  $\sigma_{ij}$ . Suppose that also the inverse matrix of  $\boldsymbol{\Sigma}$ ,  $\boldsymbol{\Sigma}^{-1}$  exists. If the joint probability density function of  $(X_1, \dots, X_n)$  is given by

$$f(\vec{x}) = \frac{1}{(2\pi)^{n/2}|\boldsymbol{\Sigma}|^{1/2}} \exp\left\{-\frac{1}{2}(\vec{x} - \vec{\mu})'\boldsymbol{\Sigma}^{-1}(\vec{x} - \vec{\mu})\right\} \text{ for } \vec{x} \in \mathbb{R}^n$$

where  $\vec{x} = (x_1, x_2, \dots, x_n)$ , then  $\vec{X}$  is said to have a **multivariate normal distribution**. We write  $\vec{X} \sim \text{MN}(\vec{\mu}, \boldsymbol{\Sigma})$ .

**REMARK — Aitken's Integral**

For any positive definite matrix  $\mathbf{A} \in \mathbb{R}^{n \times n}$ , we have

$$\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}\vec{x}'\mathbf{A}\vec{x}\right\} d\vec{x} = (2\pi)^{n/2}|\mathbf{A}|^{-1/2}.$$

**THEOREM 2.2**

If  $\vec{X} \sim MN(\vec{\mu}, \Sigma)$ , its moment generating function is given by

$$M_{\vec{X}}(\vec{t}') = \mathbb{E}[e^{\vec{t}'\vec{X}}] = \exp\left\{\vec{t}'\vec{\mu} + \frac{\vec{t}'\Sigma\vec{t}}{2}\right\}.$$

**Proof:**

$$\begin{aligned} M_{\vec{X}}(\vec{t}') &= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{\vec{t}'\vec{x} - \frac{1}{2}(\vec{x} - \vec{\mu})'\Sigma^{-1}(\vec{x} - \vec{\mu})\right\} d\vec{x} \\ &= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}[(\vec{x} - \vec{\mu})'\Sigma^{-1}(\vec{x} - \vec{\mu}) - 2\vec{t}'\vec{x}]\right\} d\vec{x} \\ &= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \\ &\quad \exp\left\{-\frac{1}{2}[(\vec{x}' - \vec{\mu}' - \vec{t}'\Sigma' + \vec{t}'\Sigma')\Sigma^{-1}(\vec{x} - \vec{\mu} - \Sigma\vec{t} + \Sigma\vec{t}) - 2\vec{t}'\vec{x}]\right\} d\vec{x} \\ &= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \\ &\quad \exp\left\{-\frac{1}{2}(\vec{x}' - \vec{\mu}' - \vec{t}'\Sigma' + \vec{t}'\Sigma')\Sigma^{-1}(\vec{x} - \vec{\mu} - \Sigma\vec{t})\right\} \\ &\quad \exp\left\{-\frac{1}{2}[\vec{x}'\vec{t} - \vec{\mu}'\vec{t} - \vec{t}'\Sigma'\vec{t} + \vec{t}'\vec{x} - \vec{t}'\vec{\mu} - \vec{t}'\Sigma\vec{t} + \vec{t}'\Sigma'\vec{t} - 2\vec{t}'\vec{x}]\right\} d\vec{x} \\ &= \underbrace{\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} (2\pi)^{-n/2} |\Sigma|^{-1/2} \exp\left\{-\frac{1}{2}(\vec{x} - \vec{\mu} - \Sigma\vec{t})'\Sigma^{-1}(\vec{x} - \vec{\mu} - \Sigma\vec{t})\right\} d\vec{x}}_{=1} \\ &\quad \exp\left\{\vec{t}'\vec{\mu} + \frac{\vec{t}'\Sigma\vec{t}}{2}\right\} \\ &= \exp\left\{\vec{t}'\vec{\mu} + \frac{\vec{t}'\Sigma\vec{t}}{2}\right\} \end{aligned}$$

**REMARK — Gamma Distribution**

$Y$  is said to have a **Gamma distribution** with shape  $\alpha$  and scale  $\beta$  when

$$f_Y(y) = \frac{y^{\alpha-1} e^{-y/\beta}}{\Gamma(\alpha)\beta^\alpha}, \text{ for } y > 0, \alpha > 0, \beta > 0,$$

and 0 otherwise. We write  $Y \sim \text{GAM}(\alpha, \beta)$ .

- (1)  $\mathbb{E}[Y] = \alpha\beta$ ,  $\text{Var}(Y) = \alpha\beta^2$ .
- (2)  $M_Y(t) = (1 - \beta t)^{-\alpha}$  for  $t < 1/\beta$ .

**REMARK — Chi-Squared Distribution**

$Q$  is said to have a **Chi-squared distribution** with  $n \in \mathbb{Z}^+$  degrees of freedom when  $Q \sim \text{GAM}(n/2, 2)$ . We write  $Q \sim \chi^2(n)$ .

- (1)  $\mathbb{E}[Q] = k$ ,  $\text{Var}(Q) = 2k$ .
- (2)  $M_Q(t) = (1 - 2t)^{-n/2}$  for  $t < 1/2$ .

**THEOREM 2.3**

Let  $\vec{X} \sim MN(\vec{0}, \Sigma)$ . Then  $\vec{X}'\Sigma^{-1}\vec{X} \sim \chi^2(n)$ .

**Proof:** Let  $Y = \vec{X}'\Sigma^{-1}\vec{X}$ . Then,

$$\begin{aligned}
M_Y(t) &= \mathbb{E}[e^{tY}] \\
&= \mathbb{E}[e^{\vec{X}'(t\Sigma^{-1})\vec{X}}] \\
&= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{\vec{x}'(t\Sigma^{-1})\vec{x} - \frac{1}{2}\vec{x}'\Sigma^{-1}\vec{x}\right\} d\vec{x} \\
&= (2\pi)^{-n/2} |\Sigma|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}\vec{x}'(1-2t)\Sigma^{-1}\vec{x}\right\} d\vec{x} \\
&= |\Sigma|^{-1/2} \left|((1-2t)\Sigma^{-1})^{-1}\right|^{1/2} \\
&= (1-2t)^{-n/2}
\end{aligned}$$

**REMARK — Non-Central Chi-Squared Distribution**

Let  $X_1, \dots, X_n$  be independent and  $X_i \sim \mathcal{N}(\mu_i, 1)$ . Set  $\lambda = \frac{1}{2}\vec{\mu}'\vec{\mu} = \frac{1}{2}\sum_{i=1}^n \mu_i^2$  and  $W = \sum_{i=1}^n X_i^2$ . Then,  $W$  has a **non-central chi-squared distribution** with degrees of freedom  $n$  and non-centrality parameter  $\lambda$ . We write  $W \sim \chi^2(n, \lambda)$ . The usual chi-square corresponds to  $\lambda = 0$ . Note: The factor  $\frac{1}{2}$  is used for this course.

**Not covered in notes:**

$$M_W(t) = (1-2t)^{-n/2} \exp\left\{\frac{\lambda 2t}{1-2t}\right\}.$$

**REMARK — F-Distribution**

If  $X \sim \chi^2(n)$  independently of  $Y \sim \chi^2(m)$  for  $n, m > 0$ , then we say  $U = \frac{X/n}{Y/m}$  has a (central) **F-distribution**. We write  $U \sim F(n, m)$ .

If  $X \sim \chi^2(n, \lambda)$  independently of  $Y \sim \chi^2(m)$  for  $n, m, \lambda > 0$ , then we say  $U = \frac{X/n}{Y/m}$  has a **non-central F-distribution**. We write  $U \sim F(n, m, \lambda)$ . If  $\lambda = 0$ , then  $U \sim F(n, m)$ .

**Transformation of Multivariate Normal**

If  $\Sigma$  is symmetric, then by the spectral theorem, there exists an orthogonal matrix  $\Gamma$  such that

$$\Sigma = \Gamma' \text{diag}(\lambda_1, \dots, \lambda_n) \Gamma,$$

where  $\lambda_1, \dots, \lambda_n$  are the eigenvalues of  $\Sigma$ . Note that  $\lambda_i > 0$  for all  $i \in [1, n]$  since  $\Sigma$  is positive definite. Furthermore, if we set

$$\Sigma^{1/2} = \Gamma' \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n}) \Gamma,$$

we see that

$$\begin{aligned}
\Sigma^{1/2}\Sigma^{1/2} &= \Gamma' \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n}) \underbrace{\Gamma\Gamma'}_{\mathbf{I}} \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n}) \Gamma \\
&= \Gamma' \text{diag}(\lambda_1, \dots, \lambda_n) \Gamma \\
&= \Sigma.
\end{aligned}$$

Therefore,  $\Sigma^{1/2}$  is well-defined and is called the **square root** of  $\Sigma$ .

**REMARK**

If  $\vec{X} \sim \text{MN}(\vec{\mu}, \Sigma)$ , and  $\vec{Z} = \Sigma^{-1/2}(\vec{X} - \vec{\mu})$ , where  $\Sigma^{-1/2} = (\Sigma^{1/2})^{-1}$ , then

$$\vec{Z} \sim \text{MN}(\vec{0}, \mathbf{I}).$$

**Proof:**

$$\begin{aligned} M_{\vec{Z}}(\vec{t}') &= \mathbb{E}[e^{\vec{t}'\vec{Z}}] \\ &= \mathbb{E}\left[\exp\{\vec{t}'\Sigma^{-1/2}(\vec{X} - \vec{\mu})\}\right] \\ &= \exp\{-\vec{t}'\Sigma^{-1/2}\vec{\mu}\} \mathbb{E}\left[\exp\{\vec{t}'\Sigma^{-1/2}\vec{X}\}\right] \\ &= \exp\{-\vec{t}'\Sigma^{-1/2}\vec{\mu}\} \exp\left\{\vec{t}'\Sigma^{-1/2}\vec{\mu} + \frac{\vec{t}'\Sigma^{-1/2}\Sigma\Sigma^{-1/2}\vec{t}}{2}\right\} \\ &= \exp\left\{\frac{\vec{t}'\Sigma^{-1/2}\Sigma^{1/2}\Sigma^{1/2}\Sigma^{-1/2}\vec{t}}{2}\right\} \\ &= \exp\left\{\frac{\vec{t}'\vec{t}}{2}\right\} \\ &= \exp\left\{\frac{1}{2}\sum_{i=1}^n t_i^2\right\}. \end{aligned}$$

### 3 Lecture 3: Some Basic Lemmas

**LEMMA 3.1**

Let  $\vec{b}$  be a vector and  $\mathbf{W}$  be a positive definite symmetric matrix. Then,

$$\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}\vec{x}'\mathbf{W}^{-1}\vec{x} + \vec{b}'\vec{x}\right\} d\vec{x} = (2\pi)^{n/2} |\mathbf{W}|^{1/2} \exp\left\{\frac{\vec{b}'\mathbf{W}\vec{b}}{2}\right\}.$$

**LEMMA 3.2**

Let  $\mathbf{A}$  be a symmetric matrix and  $\vec{X} \sim \text{MN}(\vec{\mu}, \Sigma)$ . Then,

$$M_{\vec{X}'\mathbf{A}\vec{X}}(t) = |\mathbf{I} - 2t\mathbf{A}\Sigma|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}]\Sigma^{-1}\vec{\mu}\right\}$$

for small enough  $t$ .

**Proof:** By definition,

$$\begin{aligned}
M_{\vec{X}'\mathbf{A}\vec{X}}(t) &= \mathbb{E}[\exp\{t\vec{x}'\mathbf{A}\vec{x}\}] \\
&= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \frac{1}{(2\pi)^{n/2}|\boldsymbol{\Sigma}|^{1/2}} \exp\{t\vec{x}'\mathbf{A}\vec{x}\} \exp\left\{-\frac{1}{2}(\vec{x}-\vec{\mu})'\boldsymbol{\Sigma}^{-1}(\vec{x}-\vec{\mu})\right\} d\vec{x} \\
&= (2\pi)^{-n/2}|\boldsymbol{\Sigma}|^{-1/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{\vec{x}'t\mathbf{A}\vec{x} - \frac{1}{2}(\vec{x}'\boldsymbol{\Sigma}^{-1}\vec{x} - 2\vec{x}'\boldsymbol{\Sigma}^{-1}\vec{\mu} + \vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu})\right\} d\vec{x} \\
&= (2\pi)^{-n/2}|\boldsymbol{\Sigma}|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp\left\{-\frac{1}{2}\vec{x}'\underbrace{(\mathbf{I}-2t\mathbf{A}\boldsymbol{\Sigma})\boldsymbol{\Sigma}^{-1}}_{\mathbf{W}}\vec{x} + \underbrace{\vec{\mu}'\boldsymbol{\Sigma}^{-1}}_{\vec{b}}\vec{x}\right\} d\vec{x} \\
&= (2\pi)^{-n/2}|\boldsymbol{\Sigma}|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} (2\pi)^{n/2}|\mathbf{W}^{-1}|^{1/2} \exp\left\{\frac{\vec{b}'\mathbf{W}^{-1}\vec{b}}{2}\right\} \text{ by Lemma 3.1} \\
&= |\boldsymbol{\Sigma}|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} |\mathbf{W}^{-1}|^{1/2} \exp\left\{\frac{\vec{b}'\mathbf{W}^{-1}\vec{b}}{2}\right\}.
\end{aligned}$$

Note that

$$\mathbf{W}^{-1} = [(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})\boldsymbol{\Sigma}^{-1}]^{-1} = \boldsymbol{\Sigma}(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1},$$

and

$$\begin{aligned}
\vec{b}'\mathbf{W}^{-1}\vec{b} &= \vec{\mu}'\boldsymbol{\Sigma}^{-1}\boldsymbol{\Sigma}(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1}\boldsymbol{\Sigma}^{-1}\vec{\mu} \\
&= \vec{\mu}'(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1}\boldsymbol{\Sigma}^{-1}\vec{\mu},
\end{aligned}$$

and

$$|\mathbf{W}^{-1}|^{1/2} = \left|[(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})\boldsymbol{\Sigma}^{-1}]^{-1}\right|^{1/2} = |\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma}|^{-1/2}|\boldsymbol{\Sigma}|^{1/2}.$$

Continuing,

$$\begin{aligned}
M_{\vec{X}'\mathbf{A}\vec{X}}(t) &= |\boldsymbol{\Sigma}|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} |\mathbf{W}^{-1}|^{1/2} \exp\left\{\frac{\vec{b}'\mathbf{W}^{-1}\vec{b}}{2}\right\} \\
&= \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} \exp\left\{\frac{1}{2}\vec{\mu}'(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} |\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma}|^{-1/2} \\
&= \exp\left\{-\frac{1}{2}\vec{\mu}'\boldsymbol{\Sigma}^{-1}\vec{\mu} + \frac{1}{2}\vec{\mu}'(\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\} |\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma}|^{-1/2} \\
&= \frac{1}{\sqrt{|\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma}|}} \exp\left\{-\frac{1}{2}\vec{\mu}'[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma})^{-1}]\boldsymbol{\Sigma}^{-1}\vec{\mu}\right\}.
\end{aligned}$$

### LEMMA 3.3

Let  $\lambda_1, \dots, \lambda_n$  be the eigenvalues of  $\mathbf{A}\boldsymbol{\Sigma}$ . Then,

$$|\mathbf{I} - 2t\mathbf{A}\boldsymbol{\Sigma}| = (1 - 2t\lambda_1) \cdots (1 - 2t\lambda_n).$$

**Proof:** By the spectral theorem,

$$\mathbf{A}\boldsymbol{\Sigma} = \mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n)\mathbf{Q}.$$

Then,

$$\begin{aligned}\mathbf{I} - 2t\mathbf{A}\Sigma &= \mathbf{I} - 2t\mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n)\mathbf{Q} \\ &= \mathbf{I} - \mathbf{Q}' \text{diag}(2t\lambda_1, \dots, 2t\lambda_n)\mathbf{Q} \\ &= \mathbf{Q}' \text{diag}(1 - 2t\lambda_1, \dots, 1 - 2t\lambda_n)\mathbf{Q}.\end{aligned}$$

Therefore,

$$\begin{aligned}|\mathbf{I} - 2t\mathbf{A}\Sigma| &= |\mathbf{Q}'| |\text{diag}(1 - 2t\lambda_1, \dots, 1 - 2t\lambda_n)| |\mathbf{Q}| \\ &= (1 - 2t\lambda_1) \cdots (1 - 2t\lambda_n).\end{aligned}$$

#### LEMMA 3.4

For  $t$  small enough, we have

$$\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1} = - \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r.$$

**Proof:** If  $t$  is small enough, then  $\mathbf{I} - 2t\mathbf{A}\Sigma$  is invertible. Thus,

$$(\mathbf{I} - 2t\mathbf{A}\Sigma)[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}] = \mathbf{I} - 2t\mathbf{A}\Sigma - \mathbf{I} = -2t\mathbf{A}\Sigma.$$

On the other hand,

$$\begin{aligned}(\mathbf{I} - 2t\mathbf{A}\Sigma) \left( - \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r \right) &= - \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r + \underbrace{\sum_{r=2}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r}_{\sum_{r=1}^{\infty} (2t)^{r+1} (\mathbf{A}\Sigma)^{r+1}} \\ &= -(2t)\mathbf{A}\Sigma.\end{aligned}$$

Therefore,

$$(\mathbf{I} - 2t\mathbf{A}\Sigma)[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}] = (\mathbf{I} - 2t\mathbf{A}\Sigma) \left( - \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r \right)$$

For small enough  $t$ , the inverse of  $\mathbf{I} - 2t\mathbf{A}\Sigma$  exists, so

$$\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1} = - \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r.$$

#### DEFINITION 3.1: Cumulant Generating Function

Let  $M_X(t)$  be the moment generating function of  $X$ . Then,

$$K_X(t) = \log(M_X(t))$$

is called the **cumulant generating function**. By Taylor's expansion,

$$K_X(t) = \sum_{n=1}^{\infty} \kappa_n \frac{t^n}{n!}.$$

$\kappa_n = K^{(n)}(0)$  is the  $n$ -th cumulant.

**EXAMPLE 3.1**

Let  $X \sim \mathcal{N}(\mu, \sigma^2)$ . Then,

$$M_X(t) = \exp\left\{\mu t + \frac{\sigma^2}{2}t^2\right\} \implies K_X(t) = \mu t + \frac{\sigma^2}{2}t^2 \implies \kappa_1 = \mu, \quad \kappa_2 = \sigma^2, \quad \kappa_i = 0, \quad i \geq 3.$$

**LEMMA 3.5**

For any  $X$  with  $K_X(t)$  well-defined,

$$\kappa_1 = \mathbb{E}[X], \quad \kappa_2 = \text{Var}(X).$$

**Proof:**

$$\begin{aligned} \left. \frac{dK_X(t)}{dt} \right|_{t=0} &= \left. \frac{d \log(M_X(t))}{dt} \right|_{t=0} \\ &= \left. \frac{M'_X(t)}{M_X(t)} \right|_{t=0} \\ &= M'_X(0) \\ &= \mathbb{E}[X] \\ &= \kappa_1. \end{aligned}$$

$$\begin{aligned} \left. \frac{dK'_X(t)}{dt} \right|_{t=0} &= \left. \frac{d M'_X(t)}{dt M_X(t)} \right|_{t=0} \\ &= \left. \frac{M''_X(t)M_X(t) - (M'_X(t))^2}{M_X^2(t)} \right|_{t=0} \\ &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 \\ &= \text{Var}(X) \\ &= \kappa_2. \end{aligned}$$

**THEOREM 3.1**

Let  $\vec{X} \sim MN(\vec{\mu}, \Sigma)$ . For any symmetric matrix  $\mathbf{A}$  and  $\lambda_1, \dots, \lambda_n$  are eigenvalues of  $\mathbf{A}\Sigma$

$$K_{\vec{X}'\mathbf{A}\vec{X}}(t) = -\frac{1}{2} \sum_{i=1}^n \log(1 - 2t\lambda_i) + \frac{1}{2} \vec{\mu}' \sum_{r=1}^{\infty} (2t)^r (\mathbf{A}\Sigma)^r \Sigma^{-1} \vec{\mu}.$$

$$\kappa_r = 2^{r-1} (r-1)! [\text{tr}((\mathbf{A}\Sigma)^r) + r \vec{\mu}' \mathbf{A} (\Sigma \mathbf{A})^{r-1} \vec{\mu}].$$

In particular,

$$\begin{aligned} \kappa_1 &= \mathbb{E}[\vec{X}'\mathbf{A}\vec{X}] = \text{tr}(\mathbf{A}\Sigma) + \vec{\mu}' \mathbf{A} \vec{\mu}. \\ \kappa_2 &= \text{Var}(\vec{X}'\mathbf{A}\vec{X}) \\ &= 2 [\text{tr}((\mathbf{A}\Sigma)^2) + 2 \vec{\mu}' \mathbf{A} \Sigma \mathbf{A} \vec{\mu}] \\ &= 2 \text{tr}((\mathbf{A}\Sigma)^2) + 4 \vec{\mu}' \mathbf{A} \Sigma \mathbf{A} \vec{\mu}. \end{aligned}$$

**Proof:**



Step 1: Since  $\vec{X} \sim \text{MN}(\vec{\mu}, \Sigma)$ , by Lemma 3.2,

$$M_{\vec{X}'\mathbf{A}\vec{X}}(t) = |\mathbf{I} - 2t\mathbf{A}\Sigma|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}]\Sigma^{-1}\vec{\mu}\right\}.$$

Therefore,

$$\begin{aligned} K_{\vec{X}'\mathbf{A}\vec{X}}(t) &= \log(M_{\vec{X}'\mathbf{A}\vec{X}}(t)) \\ &= \log\left(|\mathbf{I} - 2t\mathbf{A}\Sigma|^{-1/2} \exp\left\{-\frac{1}{2}\vec{\mu}'[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}]\Sigma^{-1}\vec{\mu}\right\}\right) \\ &= -\frac{1}{2}\log(|\mathbf{I} - 2t\mathbf{A}\Sigma|) - \frac{1}{2}\vec{\mu}'[\mathbf{I} - (\mathbf{I} - 2t\mathbf{A}\Sigma)^{-1}]\Sigma^{-1}\vec{\mu} \\ &= -\frac{1}{2}\log((1 - 2t\lambda_1)\cdots(1 - 2t\lambda_n)) - \frac{1}{2}\vec{\mu}'\left(-\sum_{r=1}^{\infty}(2t)^r(\mathbf{A}\Sigma)^r\right)\Sigma^{-1}\vec{\mu} \quad \text{Lemma 3.3, 3.4} \\ &= -\frac{1}{2}\sum_{i=1}^n \log(1 - 2t\lambda_i) + \frac{1}{2}\vec{\mu}'\sum_{r=1}^{\infty}(2t)^r(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}. \end{aligned}$$

Step 2: The Taylor expansion for  $\log(1 - x)$  is

$$-x - \frac{x^2}{2} - \frac{x^3}{3} - \cdots = -\sum_{r=1}^{\infty} \frac{x^r}{r}.$$

Therefore, using the Taylor expansion of  $\log(1 - 2t\lambda_i)$ , we may rewrite the first term of  $K_{\vec{X}'\mathbf{A}\vec{X}}(t)$  as

$$\begin{aligned} -\frac{1}{2}\sum_{i=1}^n \log(1 - 2t\lambda_i) &= \frac{1}{2}\sum_{i=1}^n \sum_{r=1}^{\infty} \frac{(2t\lambda_i)^r}{r} \\ &= \frac{1}{2}\sum_{r=1}^{\infty} \frac{(2t)^r \sum_{i=1}^n \lambda_i^r}{r} \\ &= \frac{1}{2}\sum_{r=1}^{\infty} \frac{(r-1)!(2t)^r \text{tr}((\mathbf{A}\Sigma)^r)}{r!} \quad \text{since } \frac{(r-1)!}{r!} = \frac{1}{r}. \end{aligned}$$

Step 3: Rewrite the second term of  $K_{\vec{X}'\mathbf{A}\vec{X}}(t)$  as

$$\frac{1}{2}\vec{\mu}'\sum_{r=1}^{\infty}(2t)^r(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu} = \frac{1}{2}\sum_{r=1}^{\infty}(2t)^r(\vec{\mu}'(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}) = \frac{1}{2}\sum_{r=1}^{\infty} \frac{r!2^r t^r \vec{\mu}'(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}}{r!}.$$

Step 4: Combining steps 1 to 3, we get

$$\begin{aligned} K_{\vec{X}'\mathbf{A}\vec{X}}(t) &= \frac{1}{2}\sum_{r=1}^{\infty} \frac{(r-1)!(2t)^r \text{tr}((\mathbf{A}\Sigma)^r)}{r!} + \frac{r!2^r t^r \vec{\mu}'(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}}{r!} \\ &= \sum_{r=1}^{\infty} 2^{r-1}(r-1)! \frac{[\text{tr}((\mathbf{A}\Sigma)^r) + r\vec{\mu}'(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}]t^r}{r!} \\ &= \sum_{r=1}^{\infty} 2^{r-1}(r-1)! \underbrace{[\text{tr}((\mathbf{A}\Sigma)^r) + r\vec{\mu}'(\mathbf{A}\Sigma)^r\Sigma^{-1}\vec{\mu}]}_{\kappa_r} \frac{t^r}{r!}, \end{aligned}$$

noting that  $(\mathbf{A}\Sigma)^r\Sigma^{-1} = (\Sigma\mathbf{A})^{r-1}$  to get the desired result.

## 4 Lecture 4: Quadratic Forms with Idempotency

### LEMMA 4.1

Let  $\mathbf{A} \in \mathbb{R}^{n \times n}$  be symmetric and  $\mathbf{B} \in \mathbb{R}^{n \times n}$  be positive definite. If the eigenvalues of  $\mathbf{AB}$  are 0's or 1's, then  $\mathbf{AB}$  is idempotent.

**Proof:** By Cholesky decomposition, there exists an invertible lower triangular matrix  $\mathbf{L}$  such that

$$\mathbf{B} = \mathbf{L}\mathbf{L}'.$$

If the eigenvalues of  $\mathbf{AB}$  are 0's or 1's, then the equation  $|\mathbf{AB} - \lambda\mathbf{I}| = 0$  has roots 0 or 1.

$$\begin{aligned} |\mathbf{AB} - \lambda\mathbf{I}| &= |\mathbf{L}'(\mathbf{AB} - \lambda\mathbf{I})(\mathbf{L}')^{-1}| \\ &= |\mathbf{L}'\mathbf{AB}(\mathbf{L}')^{-1} - \lambda\mathbf{I}| \\ &= |\mathbf{L}'\mathbf{ALL}'(\mathbf{L}')^{-1} - \lambda\mathbf{I}| \\ &= |\mathbf{L}'\mathbf{AL} - \lambda\mathbf{I}| \\ &= 0 \end{aligned}$$

has roots 0 or 1. Since  $\mathbf{L}'\mathbf{AL}$  is symmetric, and thus diagonalizable, it follows that

$$\mathbf{L}'\mathbf{AL}$$

is idempotent since

$$\begin{aligned} (\mathbf{L}'\mathbf{AL})(\mathbf{L}'\mathbf{AL}) &= \mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n) \underbrace{\mathbf{Q}\mathbf{Q}'}_{\mathbf{I}} \text{diag}(\lambda_1, \dots, \lambda_n) \mathbf{Q} \\ &= \mathbf{Q}' \text{diag}(\lambda_1^2, \dots, \lambda_n^2) \mathbf{Q} \\ &= \mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n) \mathbf{Q} \\ &= \mathbf{L}'\mathbf{AL}. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbf{L}'\mathbf{AL} &= \mathbf{L}'\mathbf{ALL}'\mathbf{AL} \\ \implies \mathbf{AL} &= \mathbf{ALL}'\mathbf{AL} \\ \implies \mathbf{ALL}' &= \mathbf{AB} = \mathbf{ALL}'\mathbf{ALL}' = \mathbf{ABAB} \\ \implies \mathbf{AB} &\text{ is idempotent.} \end{aligned}$$

### THEOREM 4.1

Let  $\vec{X} \sim MN(\vec{\mu}, \Sigma)$  and  $\mathbf{A}$  be a symmetric matrix with rank  $r$ . Then,

$$\vec{X}'\mathbf{A}\vec{X} \sim \chi^2(r, \lambda), \quad \forall \vec{\mu}$$

with  $\lambda = \frac{1}{2} \vec{\mu}'\mathbf{A}\vec{\mu}$  if and only if  $\mathbf{A}\Sigma$  is idempotent.

**Proof:**

( $\Leftarrow$ ) Assume that  $\mathbf{A}\Sigma$  is idempotent, then all eigenvalues of  $\mathbf{A}\Sigma$  are 1 or 0 (which we denote as  $\lambda_i$ ).

Since  $\Sigma$  has full rank,

$$\text{rank}(\mathbf{A}\Sigma) = \text{rank}(\mathbf{A}) = r.$$

Therefore,  $r$  eigenvalues are 1 and  $n - r$  are 0. By Theorem 3.1, we have

$$\begin{aligned} M_{\vec{X}', \mathbf{A}\vec{X}}(t) &= \prod_{i=1}^n (1 - 2t\lambda_i)^{-1/2} \exp \left\{ \frac{1}{2} \vec{\mu}' \sum_{j=1}^{\infty} (2t)^j (\mathbf{A}\Sigma)^j \Sigma^{-1} \vec{\mu} \right\} \\ &= (1 - 2t)^{-r/2} \exp \left\{ \frac{1}{2} \vec{\mu}' \sum_{j=1}^{\infty} (2t)^j \mathbf{A}\Sigma \Sigma^{-1} \vec{\mu} \right\} \\ &= (1 - 2t)^{-r/2} \exp \left\{ \frac{1}{2} \vec{\mu}' \mathbf{A} \vec{\mu} \sum_{j=1}^{\infty} (2t)^j \right\} \\ &= (1 - 2t)^{-r/2} \exp \left\{ \frac{\vec{\mu}' \mathbf{A} \vec{\mu}}{2} \frac{2t}{1 - 2t} \right\} \\ &= (1 - 2t)^{-r/2} \exp \left\{ \lambda \frac{2t}{1 - 2t} \right\}. \end{aligned}$$

Let  $\eta \sim \chi^2(r, \lambda)$ . By definition,

$$\eta = X_1^2 + \dots + X_r^2,$$

where  $X_i \sim \mathcal{N}(\mu_i, 1)$  and  $X_1, \dots, X_r$  are independent.

$$\begin{aligned} M_{\eta}(t) &= \mathbb{E}[e^{t\eta}] \\ &= \mathbb{E}[e^{t(X_1^2 + \dots + X_r^2)}] \\ &= \prod_{i=1}^r \mathbb{E}[e^{tX_i^2}]. \end{aligned}$$

Now,

$$\begin{aligned} \mathbb{E}[e^{tX_i^2}] &= \int_{-\infty}^{\infty} (2\pi)^{-1/2} \exp\{tx^2\} \exp\left\{-\frac{(x - \mu_i)^2}{2}\right\} dx \\ &= \int_{-\infty}^{\infty} (2\pi)^{-1/2} \exp\left\{-\frac{1}{2}(x^2 - 2tx^2 - 2\mu_i x + \mu_i^2)\right\} dx \\ &= (1 - 2t)^{-1/2} \exp\left\{-\frac{1 - 2t}{2} \left(\frac{\mu_i^2}{1 - 2t} - \left(\frac{\mu_i}{1 - 2t}\right)^2\right)\right\} \\ &\quad \underbrace{\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}(1 - 2t)^{-1/2}} \exp\left\{-\frac{(x - \frac{\mu_i}{1 - 2t})^2}{2(1 - 2t)^{-1}}\right\} dx}_{\mathcal{N}\left(\frac{\mu_i}{1 - 2t}, (1 - 2t)^{-1}\right)} \\ &= (1 - 2t)^{-1/2} \exp\left\{\mu_i^2 \frac{t}{1 - 2t}\right\} \\ &= (1 - 2t)^{-1/2} \exp\left\{\frac{\mu_i^2}{2} \frac{2t}{1 - 2t}\right\}. \end{aligned}$$

Hence,

$$\begin{aligned}
M_{\eta}(t) &= \prod_{i=1}^r (1-2t)^{-1/2} \exp\left\{\frac{\mu_i^2}{2} \frac{2t}{1-2t}\right\} \\
&= (1-2t)^{-r/2} \exp\left\{\frac{1}{2} \underbrace{\sum_{i=1}^n \mu_i^2}_{\lambda} \frac{2t}{1-2t}\right\} \\
&= (1-2t)^{-r/2} \exp\left\{\frac{\lambda 2t}{1-2t}\right\},
\end{aligned}$$

which is the mgf of  $\chi^2(r, \lambda)$ . By uniqueness of moment generating functions,

$$\vec{X}' \mathbf{A} \vec{X} \sim \chi^2(r, \lambda).$$

( $\implies$ ) Assume  $\vec{X}' \mathbf{A} \vec{X} \sim \chi^2(r, \lambda)$  for all  $\vec{\mu}$ . Choose  $\vec{\mu} = \vec{0}$ , then  $\lambda = 0$ .

$$\begin{aligned}
M_{\vec{X}' \mathbf{A} \vec{X}}(t) &= \prod_{i=1}^n (1-2t\lambda_i)^{-1/2} \\
&= (1-2t)^{-r/2}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\implies \prod_{i=1}^n (1-2t\lambda_i) &= (1-2t)^r && \text{cancel exponents} \\
\implies \sum_{i=1}^n \log(1-2t\lambda_i) &= r \log(1-2t) && \text{take logarithm} \\
\implies \sum_{i=1}^n \left[ \sum_{\ell=1}^{\infty} \frac{(2t\lambda_i)^\ell}{\ell} \right] &= r \sum_{\ell=1}^{\infty} \frac{(2t)^\ell}{\ell} && \text{Taylor expansion} \\
\implies \sum_{\ell=1}^{\infty} \frac{(\sum_{i=1}^n \lambda_i^\ell - r)(2t)^\ell}{\ell} &= 0 && \text{re-order summation}
\end{aligned}$$

Therefore,

$$\sum_{i=1}^n \lambda_i^\ell = r, \quad \forall \ell \geq 1.$$

Case 1: If  $|\lambda_i| \geq 1$  for some  $i$ , then choose  $\ell = 2k$  and let  $2k \rightarrow \infty$ , then

$$\lambda_1^{2k} + \dots + \lambda_i^{2k} + \dots + \lambda_n^{2k} = r,$$

but the left-hand side is  $\infty \neq r$ , contradiction. Thus,  $|\lambda_i| \leq 1$  for all  $i$ .

Case 2: If  $|\lambda_i| < 1$  for some  $i$ , then choose  $\ell = 2k$  and let  $k \rightarrow \infty$ ,  $|\lambda_i|^{2k} \rightarrow 0$ . Hence, the total terms with  $|\lambda_i| < 1$  will be  $n - r$ . The equality

$$\lambda_1 + \dots + \lambda_n = r$$

implies that all the terms with  $|\lambda_i| = 1$  are actually  $\lambda_i$ . Why? Let  $d$  be the number of  $i$  such that  $|\lambda_i| < 1$ , there will be  $n - d$  of  $|\lambda_i| = 1$  (fill in details).

Hence, the eigenvalues of  $\mathbf{A}\Sigma$  are 1 or 0. Since  $\Sigma$  is positive definite, it follows from Lemma 4.1 that  $\mathbf{A}\Sigma$  is idempotent.

**EXAMPLE 4.1**

Let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$ . Then,

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1).$$

**Solution:** Let

$$\vec{X} = \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}.$$

Then,

$$\vec{X} \sim \text{MN}(\mu \vec{j}, \sigma^2 \mathbf{I}).$$

$$\begin{aligned} \frac{(n-1)S^2}{\sigma^2} &= \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \bar{X})^2 \\ &= \frac{1}{\sigma^2} \vec{X}' (\mathbf{I} - \frac{1}{n} \mathbf{J}) \vec{X} \\ &= \vec{Y}' \mathbf{A} \vec{Y}, \end{aligned}$$

where  $\vec{Y} = \frac{1}{\sigma} \vec{X}$  and  $\mathbf{A} = \mathbf{I} - \frac{1}{n} \mathbf{J}$ , so

$$\vec{Y} \sim \text{MN}\left(\frac{\mu}{\sigma} \vec{j}, \mathbf{I}\right).$$

$$\mathbf{A} \Sigma = (\mathbf{I} - \frac{1}{n} \mathbf{J}) \mathbf{I} = \mathbf{I} - \frac{1}{n} \mathbf{J} = \mathbf{A}.$$

Also,

$$\mathbf{A}^2 = \mathbf{I} - \frac{2}{n} \mathbf{J} + \frac{1}{n^2} \mathbf{J} \mathbf{J} = \mathbf{I} - \frac{1}{n} \mathbf{J} = \mathbf{A}.$$

Therefore,  $\mathbf{A} \Sigma$  is idempotent. By Theorem 5.1,

$$\vec{Y}' \mathbf{A} \vec{Y} = \frac{(n-1)S^2}{\sigma^2} \sim \chi^2(r, \lambda),$$

with

$$r = \text{rank}(\mathbf{A}) = \text{rank}(\mathbf{I} - \frac{1}{n} \mathbf{J}) = n - 1$$

and

$$\begin{aligned} \lambda &= \frac{1}{2} \vec{\mu}' \mathbf{A} \vec{\mu} \\ &= \frac{1}{2} \frac{\mu}{\sigma} \vec{j}' (\mathbf{I} - \frac{1}{n} \mathbf{J}) \frac{\mu}{\sigma} \vec{j} \\ &= \frac{\mu^2}{2\sigma^2} (\vec{j}' \vec{j} - \frac{1}{n} \vec{j}' \mathbf{J} \vec{j}) \\ &= \frac{\mu^2}{2\sigma^2} (n - \frac{1}{n} n^2) \\ &= 0. \end{aligned}$$

Therefore,  $\vec{Y}' \mathbf{A} \vec{Y} \sim \chi^2(r, 0) = \chi^2(r)$ .

## 5 Lecture 5: Criteria for Independence

### LEMMA 5.1

Let  $\mathbf{A}$  be a symmetric positive semidefinite  $n \times n$  matrix with rank  $r$ . Then, there exists an  $n \times r$  matrix  $\mathbf{D}$  with rank  $r$  such that

$$\mathbf{A} = \mathbf{D}\mathbf{D}'$$

**Proof:** If  $\mathbf{A}$  is symmetric, then by the spectral theorem

$$\mathbf{A} = \mathbf{Q}' \text{diag}(\lambda_1, \dots, \lambda_n) \mathbf{Q},$$

where  $\mathbf{Q}$  is orthogonal where  $r$  of  $\lambda_i$  are non-zero since  $\text{rank}(\mathbf{A}) = r$ . Without loss of generality, we can assume that  $\lambda_1, \dots, \lambda_r > 0, \lambda_j = 0$  for  $j > r$ . Define

$$\mathbf{D} = \mathbf{Q}' \begin{bmatrix} \mathbf{\Lambda}^{1/2} \\ \mathbf{O} \end{bmatrix}_{n \times r},$$

where

$$\mathbf{\Lambda}^{1/2} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_r}).$$

Hence,

$$\mathbf{D}\mathbf{D}' = \mathbf{Q}' \begin{bmatrix} \mathbf{\Lambda}^{1/2} \\ \mathbf{O} \end{bmatrix} \begin{bmatrix} \mathbf{\Lambda}^{1/2} & \mathbf{O} \end{bmatrix} \mathbf{Q} = \mathbf{Q}' \begin{bmatrix} \mathbf{\Lambda} & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{bmatrix} \mathbf{Q} = \mathbf{A}.$$

### THEOREM 5.1

Let  $\vec{X} \sim MN(\vec{\mu}, \Sigma)$ ,  $\mathbf{A} \in \mathbb{R}^{n \times n}$  be symmetric and  $\mathbf{B} \in \mathbb{R}^{k \times n}$ .  $\vec{X}'\mathbf{A}\vec{X}$  and  $\mathbf{B}\vec{X}$  are independent if and only if  $\mathbf{B}\Sigma\mathbf{A} = \mathbf{O}$ .

**Proof:** We assume that  $\mathbf{A}$  is positive semidefinite.

Step 1: Let  $r = \text{rank}(\mathbf{A})$ ,  $\mathbf{A} = \mathbf{D}\mathbf{D}'$  from Lemma 5.1. We know that  $\mathbf{D}$  is  $n \times r$  with  $\text{rank}(\mathbf{D}) = r$ , and since  $\text{rank}(\mathbf{D}\mathbf{D}') = \text{rank}(\mathbf{D}) = r$ , then  $\mathbf{D}'\mathbf{D}$  is invertible. We will show that

$$\mathbf{B}\Sigma\mathbf{A} = \mathbf{O} \iff \mathbf{B}\Sigma\mathbf{D} = \mathbf{O}.$$

( $\implies$ ) Note that

$$\begin{aligned} \mathbf{B}\Sigma\mathbf{A} &= \mathbf{B}\Sigma\mathbf{D}\mathbf{D}' = \mathbf{O} \\ &\implies \mathbf{B}\Sigma\mathbf{D}\mathbf{D}'\mathbf{D} = \mathbf{O} \\ &\implies \mathbf{B}\Sigma\mathbf{D}\mathbf{D}'\mathbf{D}(\mathbf{D}'\mathbf{D})^{-1} = \mathbf{B}\Sigma\mathbf{D} = \mathbf{O}. \end{aligned}$$

On the other hand, if  $\mathbf{B}\Sigma\mathbf{D} = \mathbf{O}$ , then

$$\mathbf{B}\Sigma\mathbf{D}\mathbf{D}' = \mathbf{B}\Sigma\mathbf{A} = \mathbf{O}.$$

Step 2 (Sufficiency): Assume that  $\mathbf{B}\Sigma\mathbf{A} = \mathbf{O}$ , then  $\mathbf{B}\Sigma\mathbf{D} = \mathbf{O}$ . By direct calculation,

$$\text{Cov}(\mathbf{B}\vec{X}, \vec{X}'\mathbf{D}) = \mathbf{B} \text{Cov}(\vec{X}, \vec{X})\mathbf{D} = \mathbf{B}\Sigma\mathbf{D} = \mathbf{O}.$$

Since  $\mathbf{B}\vec{X}$  and  $\vec{X}'\mathbf{D}$  are multivariate normal, it follows that  $\mathbf{B}\vec{X}$  and  $\vec{X}'\mathbf{D}$  are independent. Noting that

$$\begin{aligned} \vec{X}'\mathbf{A}\vec{X} &= \vec{X}'\mathbf{D}\mathbf{D}'\vec{X} && \text{Lemma 5.1} \\ &= (\vec{X}'\mathbf{D})(\vec{X}'\mathbf{D})', \end{aligned}$$

which is a function of  $\vec{X}'\mathbf{D}$ . We know that if  $X$  and  $Y$  are independent, then for any measurable function  $f(X)$  and  $g(Y)$  are independent. Hence,  $\mathbf{B}\vec{X}$  and  $\vec{X}'\mathbf{A}\vec{X}$  are independent.

Step 3 (Necessity): Assume that  $\mathbf{B}\vec{X}$  and  $\vec{X}'\mathbf{A}\vec{X}$  are independent. By direct calculation,

$$\begin{aligned}
\text{Cov}(\mathbf{B}\vec{X}, \vec{X}'\mathbf{A}\vec{X}) &= \mathbf{B} \text{Cov}(\vec{X}, \vec{X}'\mathbf{A}\vec{X}) \\
&= \mathbf{B} \mathbb{E}[(\vec{X} - \vec{\mu})(\vec{X}'\mathbf{A}\vec{X} - \mathbb{E}[\vec{X}'\mathbf{A}\vec{X}])] \\
&= \mathbf{B} \mathbb{E}[(\vec{X} - \vec{\mu})(\vec{X}'\mathbf{A}\vec{X} - \vec{\mu}'\mathbf{A}\vec{\mu} - \text{tr}(\mathbf{A}\Sigma))] \\
&= \mathbf{B} \mathbb{E}[(\vec{X} - \vec{\mu})(\vec{X}'\mathbf{A}\vec{X} - \vec{\mu}'\mathbf{A}\vec{\mu})] + \underbrace{\mathbf{B} \mathbb{E}[(\vec{X} - \vec{\mu})]}_0 \text{tr}(\mathbf{A}\Sigma) \\
&= \mathbf{B} \mathbb{E}[(\vec{X} - \vec{\mu})(\vec{X}'\mathbf{A}\vec{X} - \vec{\mu}'\mathbf{A}\vec{\mu})] \\
&= \mathbf{B} \mathbb{E}\left[(\vec{X} - \vec{\mu})[(\vec{X} - \vec{\mu})'(\vec{X} - \vec{\mu}) + 2(\vec{X} - \vec{\mu})'\mathbf{A}\vec{\mu}]\right] \\
&= \mathbf{B} \mathbb{E}\left[(\vec{X} - \vec{\mu})(\vec{X} - \vec{\mu})'\mathbf{A}(\vec{X} - \vec{\mu})\right] + 2\mathbf{B} \mathbb{E}\left[(\vec{X} - \vec{\mu})'\mathbf{A}\vec{\mu}\right] \\
&= \mathbf{B} \mathbb{E}\left[(\vec{X} - \vec{\mu})(\vec{X} - \vec{\mu})'\mathbf{A}(\vec{X} - \vec{\mu})\right] + 2\mathbf{B}\Sigma\mathbf{A}\vec{\mu}.
\end{aligned}$$

To show that the first term is zero, using the spectral theorem re-write  $\mathbf{A}$ , define  $\vec{Y} = \vec{X} - \vec{\mu}$ , and use the fact that the third moments of multivariate normal are 0 (exercise). Hence,

$$\mathbf{B}\Sigma\mathbf{A}\vec{\mu} = \mathbf{0}.$$

Since  $\vec{\mu}$  is arbitrary, it follows that

$$\mathbf{B}\Sigma\mathbf{A} = \mathbf{0}.$$

### THEOREM 5.2

Let  $\vec{X} \sim MN(\vec{\mu}, \Sigma)$ ,  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$  be symmetric matrices.  $\vec{X}'\mathbf{A}\vec{X}$  and  $\vec{X}'\mathbf{B}\vec{X}$  are independent if and only if  $\mathbf{A}\Sigma\mathbf{B} = \mathbf{0}$ .

**Proof:** Let  $\text{rank}(\mathbf{A}) = r$ ,  $\text{rank}(\mathbf{B}) = s$ . By the spectral theorem, there are orthogonal matrices  $\mathbf{Q}_1$  and  $\mathbf{Q}_2$  such that

$$\begin{aligned}
\mathbf{A} &= \mathbf{Q}'_1 \text{diag}(\lambda_1, \dots, \lambda_n) \mathbf{Q}_1, \\
\mathbf{B} &= \mathbf{Q}'_2 \text{diag}(\tilde{\lambda}_1, \dots, \tilde{\lambda}_n) \mathbf{Q}_2.
\end{aligned}$$

Without loss of generality, we assume that

- $\lambda_1, \dots, \lambda_r \neq 0$ ,  $\lambda_j = 0$  for  $j > r$ ,
- $\tilde{\lambda}_1, \dots, \tilde{\lambda}_s \neq 0$ ,  $\tilde{\lambda}_i = 0$  for  $i > s$ .

Set

$$\mathbf{D}_r = \text{diag}(\lambda_1, \dots, \lambda_r), \quad \tilde{\mathbf{D}}_s = \text{diag}(\tilde{\lambda}_1, \dots, \tilde{\lambda}_s).$$

Hence,  $\mathbf{Q}'_1 = (\mathbf{Q}'_{11} \quad \mathbf{Q}'_{12})$  with  $\mathbf{Q}'_{11}$  being  $n \times r$  and  $\text{rank}(\mathbf{Q}'_{11}) = r$ . Then,

$$\begin{aligned}
\mathbf{A} &= (\mathbf{Q}'_{11} \quad \mathbf{Q}'_{12}) \begin{pmatrix} \mathbf{D}_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{Q}_{11} \\ \mathbf{Q}_{12} \end{pmatrix} \\
&= \mathbf{Q}'_{11} \mathbf{D}_r \mathbf{Q}_{12}.
\end{aligned}$$

Define  $\mathbf{Q}'_2 = (\tilde{\mathbf{Q}}'_{11} \quad \tilde{\mathbf{Q}}'_{12})$  to similarly get

$$\mathbf{B} = \tilde{\mathbf{Q}}'_{11} \tilde{\mathbf{D}}_s \tilde{\mathbf{Q}}_{12}.$$

( $\Leftarrow$ ) “Sufficiency:” Assume that  $\mathbf{A}\Sigma\mathbf{B} = \mathbf{O}$ , so

$$\begin{aligned}\mathbf{A}\Sigma\mathbf{B} &= \mathbf{Q}'_{11}\mathbf{D}_r\mathbf{Q}_{11}\Sigma\tilde{\mathbf{Q}}'_{11}\tilde{\mathbf{D}}_s\tilde{\mathbf{Q}}_{11} = \mathbf{O} \\ \Rightarrow \mathbf{Q}_{11}\mathbf{Q}'_{11}\mathbf{Q}'_{11}\mathbf{D}_r\mathbf{Q}_{11}\Sigma\tilde{\mathbf{Q}}'_{11}\tilde{\mathbf{D}}_s\tilde{\mathbf{Q}}_{11} &= \mathbf{O} \\ \Rightarrow \mathbf{D}_r\mathbf{Q}_{11}\Sigma\tilde{\mathbf{Q}}'_{11}\tilde{\mathbf{D}}_s &= \mathbf{O} \\ \Rightarrow \mathbf{Q}_{11}\Sigma\tilde{\mathbf{Q}}'_{11} &= \mathbf{O}\end{aligned}$$

Noting that

$$\text{Cov}(\mathbf{Q}_{11}\vec{X}, \vec{X}'\tilde{\mathbf{Q}}'_{11}) = \mathbf{Q}_{11}\Sigma\tilde{\mathbf{Q}}'_{11} = \mathbf{O}.$$

Therefore,  $\mathbf{Q}_{11}\vec{X}$  and  $\vec{X}'\tilde{\mathbf{Q}}'_{11}$  are independent. Hence,

$$\begin{aligned}\vec{X}'\mathbf{A}\vec{X} &= \vec{X}'\mathbf{Q}_{11}\mathbf{D}_r\mathbf{Q}'_{11}\vec{X} \\ &= (\mathbf{Q}'_{11}\vec{X})'\mathbf{D}_r\mathbf{Q}'_{11}\vec{X}\end{aligned}$$

is a function of  $\mathbf{Q}'_{11}\vec{X}$ , and similarly  $\vec{X}'\mathbf{B}\vec{X}$  is a function of  $\vec{X}'\tilde{\mathbf{Q}}'_{11}$ . Therefore,  $\vec{X}'\mathbf{A}\vec{X}$  is independent of  $\vec{X}'\mathbf{B}\vec{X}$ .

( $\Rightarrow$ ) “Necessity:” Assume that  $\vec{X}'\mathbf{A}\vec{X}$  and  $\vec{X}'\mathbf{B}\vec{X}$  are independent. By Theorem 3.1, we have

$$\begin{aligned}\text{Var}(\vec{X}'\mathbf{A}\vec{X}) &= 2\text{tr}((\mathbf{A}\Sigma)^2) + 4\vec{\mu}'\mathbf{A}\Sigma\mathbf{A}\vec{\mu}. \\ \text{Var}(\vec{X}'\mathbf{B}\vec{X}) &= 2\text{tr}((\mathbf{B}\Sigma)^2) + 4\vec{\mu}'\mathbf{B}\Sigma\mathbf{B}\vec{\mu}.\end{aligned}$$

Since  $(\mathbf{A} + \mathbf{B})$  is symmetric,

$$\begin{aligned}\text{Var}(\vec{X}'(\mathbf{A} + \mathbf{B})\vec{X}) &= \text{Var}(\vec{X}'\mathbf{A}\vec{X} + \vec{X}'\mathbf{B}\vec{X}) \\ &= \text{Var}(\vec{X}'\mathbf{A}\vec{X}) + \text{Var}(\vec{X}'\mathbf{B}\vec{X}) \quad \text{by assumption.}\end{aligned}$$

Hence,

$$2\text{tr}(((\mathbf{A} + \mathbf{B})\Sigma)^2) + 4\vec{\mu}'(\mathbf{A} + \mathbf{B})\Sigma(\mathbf{A} + \mathbf{B})\vec{\mu} = 2\text{tr}((\mathbf{A}\Sigma)^2) + 2\text{tr}((\mathbf{B}\Sigma)^2) + 4\vec{\mu}'(\mathbf{A}\Sigma\mathbf{A} + \mathbf{B}\Sigma\mathbf{B})\vec{\mu}.$$

Therefore,

$$2\text{tr}((\mathbf{A}\Sigma\mathbf{B}\Sigma) + \text{tr}(\mathbf{B}\Sigma\mathbf{A}\Sigma)) + 4\vec{\mu}'(\mathbf{A}\Sigma\mathbf{B} + \mathbf{B}\Sigma\mathbf{A})\vec{\mu} = 0.$$

By cyclic property of trace, we obtain

$$\text{tr}(\mathbf{A}\Sigma\mathbf{B}\Sigma) = \text{tr}(\Sigma\mathbf{B}\Sigma\mathbf{A}) = \text{tr}(\mathbf{B}\Sigma\mathbf{A}\Sigma).$$

On the other hand,

$$\vec{\mu}'\mathbf{A}\Sigma\mathbf{B}\vec{\mu} + 4\vec{\mu}'\mathbf{A}\Sigma\mathbf{B}\vec{\mu} = 0.$$

Choose  $\vec{\mu} = \vec{0}$ , we get

$$\text{tr}(\mathbf{A}\Sigma\mathbf{B}\Sigma) = 0.$$

Thus,  $\vec{\mu}'\mathbf{A}\Sigma\mathbf{B}\vec{\mu} = 0$  for all  $\vec{\mu}$ , which implies that

$$\mathbf{A}\Sigma\mathbf{B} = \mathbf{O}.$$

### EXAMPLE 5.1

Let  $\vec{X}' = (X_1, X_2)' \sim \text{MN}(\vec{\mu}, \mathbf{I}_2)$ . Show

$$(X_1 - X_2)^2 \text{ is independent of } (X_1 + X_2)^2.$$



**Solution:**

$$(X_1 - X_2)^2 = X_1^2 - 2X_1X_2 + X_2^2 = X_1^2 - X_1X - 2 - X_2X_1 + X_2^2 = (X_1 \ X_2) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}.$$

$$(X_1 + X_2)^2 = X_1^2 + X_1X_2 + X_2X_1 + X_2^2 = (X_1 \ X_2) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}.$$

Now,

$$\mathbf{A}\Sigma\mathbf{B} = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \mathbf{O},$$

as required. Therefore,  $(X_1 - X_2)^2$  is independent of  $(X_1 + X_2)^2$  by Theorem 5.2.

### EXAMPLE 5.2

Let  $\vec{X}' = (X_1, X_2)' \sim MN(\vec{\mu}, \Sigma)$ . Define

$$\mathbf{B} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}.$$

Find

$$\mathbf{A} = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$$

such that  $\vec{X}'\mathbf{A}\vec{X}$  is independent of  $\vec{X}'\mathbf{B}\vec{X}$ , where

$$\Sigma = \begin{pmatrix} 1 & -1/2 \\ -1/2 & 1 \end{pmatrix}.$$

**Solution:**

$$\vec{X}'\mathbf{B}\vec{X} = X_1^2 + 4X_1X_2 + X_2^2 = (X_1 + X_2)^2 + 2X_1X_2.$$

$$\begin{aligned} \mathbf{A}\Sigma\mathbf{B} &= \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} 1 & -1/2 \\ -1/2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \\ &= \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} 0 & 1.5 \\ 1.5 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 1.5b & 1.5a \\ 1.5c & 1.5b \end{pmatrix} \\ &= 1.5 \begin{pmatrix} b & a \\ c & b \end{pmatrix} \\ &= \mathbf{O} \end{aligned}$$

implies that  $a = b = c = 0$ , so  $\mathbf{A} = \mathbf{O}$ . Therefore, there is no quadratic form.

## 6 Lecture 6: Cochran's Theorem

LECTURE 6  
26th January

**LEMMA 6.1**

Let  $\mathbf{C} = \mathbf{A} + \mathbf{B}$ . Assume that  $\mathbf{A}, \mathbf{B}$  are both  $n \times n$  symmetric. If  $\mathbf{C}^2 = \mathbf{C}$ ,  $\mathbf{A}^2 = \mathbf{A}$ ,  $\mathbf{B}$  positive semidefinite,

$$\text{rank}(\mathbf{A}) + \text{rank}(\mathbf{B}) = \text{rank}(\mathbf{C}),$$

then

$$\mathbf{AB} = \mathbf{O}.$$

**Proof:** Let  $\text{rank}(\mathbf{A}) = r$ ,  $\text{rank}(\mathbf{B}) = s$ ,  $\text{rank}(\mathbf{C}) = t = r + s$ . If  $\mathbf{C}^2 = \mathbf{C}$ , then there exists an orthogonal matrix  $\mathbf{\Gamma}$  such that

$$\mathbf{\Gamma}'\mathbf{C}\mathbf{\Gamma} = \begin{pmatrix} \mathbf{I}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix}.$$

Since  $\mathbf{C} = \mathbf{A} + \mathbf{B}$ ,

$$\mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} + \mathbf{\Gamma}'\mathbf{B}\mathbf{\Gamma} = \begin{pmatrix} \mathbf{I}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix}.$$

$\mathbf{A}$  and  $\mathbf{B}$  are positive semidefinite implies that  $\mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma}$  and  $\mathbf{\Gamma}'\mathbf{B}\mathbf{\Gamma}$  are positive semidefinite. If the element on the diagonal is zero, then the corresponding row and columns are zeros. Hence,

$$\mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} = \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix}, \quad \mathbf{\Gamma}'\mathbf{B}\mathbf{\Gamma} = \begin{pmatrix} \mathbf{H}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix}.$$

Since  $\mathbf{A}^2 = \mathbf{A}$ , we have

$$\begin{aligned} \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma}\mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} &= \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} = \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \\ \implies \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} \begin{pmatrix} \mathbf{I}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} &= \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma} + \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma}\mathbf{\Gamma}'\mathbf{B}\mathbf{\Gamma} \\ \implies \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \begin{pmatrix} \mathbf{I}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} &= \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} + \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \begin{pmatrix} \mathbf{H}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{G}_t & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} + \begin{pmatrix} \mathbf{G}_r\mathbf{H}_r & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \\ \implies \begin{pmatrix} \mathbf{G}_r\mathbf{H}_r & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} &= \mathbf{\Gamma}'\mathbf{A}\mathbf{\Gamma}\mathbf{\Gamma}'\mathbf{B}\mathbf{\Gamma} \\ &= \mathbf{\Gamma}'\mathbf{A}\mathbf{B}\mathbf{\Gamma} = \mathbf{O}. \end{aligned}$$

Therefore,  $\mathbf{AB} = \mathbf{O}$  since  $\mathbf{\Gamma}$  is orthogonal and invertible.

**THEOREM 6.1: Cochran**

Let  $\vec{X} \sim MN(\vec{0}, \mathbf{I}_n)$ ,  $\mathbf{A}_1, \dots, \mathbf{A}_m$  be symmetric  $n \times n$  matrices with  $\text{rank}(\mathbf{A}_i) = r_i$ , and  $\sum_{i=1}^m \mathbf{A}_i = \mathbf{I}_n$ .  $\vec{X}'\mathbf{A}_i\vec{X} \sim \chi^2(r_i)$  are independent if and only if  $\sum_{i=1}^m r_i = n$ .

**Proof:** ( $\Leftarrow$ ) "Sufficiency" Assume  $\sum_{i=1}^m r_i = n$ . For each  $i = 1, \dots, m$ , set

$$\mathbf{B}_i = \mathbf{I} - \mathbf{A}_i,$$

with  $\text{rank}(\mathbf{B}_i) = s_i$ . We claim that  $s_i = n - r_i$ .

$$\begin{aligned} s_i &= \text{rank}(\mathbf{I} - \mathbf{A}_i) \\ &= \text{rank}\left(\sum_{j \neq i} \mathbf{A}_j\right) \\ &\leq \sum_{j \neq i} \text{rank}(\mathbf{A}_j) \\ &= n - r_i. \end{aligned}$$

By definition,  $\mathbf{I} = \mathbf{A}_i + \mathbf{B}_i \implies \text{rank}(\mathbf{I}) = n$ . So,

$$\begin{aligned} \text{rank}(\mathbf{I}) &= n \\ &= \text{rank}(\mathbf{A}_i + \mathbf{B}_i) \\ &\leq \text{rank}(\mathbf{A}_i) + \text{rank}(\mathbf{B}_i). \end{aligned}$$

Therefore,  $\text{rank}(\mathbf{B}_i) \geq n - r_i \implies s_i = n - r_i$  for all  $i$ . Hence,

$$|\lambda \mathbf{I} - \mathbf{B}_i| = 0$$

have  $r_i$  roots being 0. Noting that

$$\begin{aligned} |\lambda \mathbf{I} - \mathbf{B}_i| &= |(\lambda - 1)\mathbf{I} - \mathbf{A}_i| \\ &= |\tilde{\lambda} \mathbf{I} - \mathbf{A}_i| \end{aligned}$$

have  $r_i$  roots being 1. Since  $\text{rank}(\mathbf{A}_i) = r_i$ , it follows that all other roots of  $\mathbf{A}_i$  are 0. Hence,

$$\mathbf{A}_i = \mathbf{A}_i \Sigma = \mathbf{A}_i \mathbf{I}$$

is idempotent by Lemma 4.1. Write

$$\mathbf{I} = \mathbf{A}_1 + (\mathbf{A}_2 + \cdots + \mathbf{A}_m).$$

Since  $\mathbf{I}^2 = \mathbf{I}$ ,  $\mathbf{A}_1^2 = \mathbf{A}_1$ ,  $\mathbf{A}_2 + \cdots + \mathbf{A}_m$  is positive semidefinite, it follows from Lemma 6.1 that

$$\mathbf{A}_1(\mathbf{A}_2 + \cdots + \mathbf{A}_m) = \mathbf{O}.$$

This implies that

$$\mathbf{I}^2 = \mathbf{A}_1 + (\mathbf{A}_2 + \cdots + \mathbf{A}_m),$$

which implies that  $\mathbf{A}_2 + \cdots + \mathbf{A}_m$  is idempotent. Applying Lemma 6.1 to

$$\mathbf{A}_2 + \cdots + \mathbf{A}_m = \mathbf{A}_2 + (\mathbf{A}_3 + \cdots + \mathbf{A}_m)$$

it follows that

$$\mathbf{A}_2(\mathbf{A}_3 + \cdots + \mathbf{A}_m) = \mathbf{O}.$$

By induction, we get

$$\mathbf{A}_{m-1} \mathbf{A}_m = \mathbf{O}.$$

By re-labeling, we get

$$\mathbf{A}_i \mathbf{A}_j = 0 \quad \forall i \neq j.$$

Since

$$\mathbf{A}_i \Sigma \mathbf{A}_j = \mathbf{A}_i \mathbf{A}_j = \mathbf{O},$$

it follows from Theorem 5.2 that

$$\vec{X}' \mathbf{A}_i \vec{X} \text{ and } \vec{X}' \mathbf{A}_j \vec{X}$$

are independent. The fact that

$$\vec{X}' \mathbf{A}_i \vec{X} \sim \chi^2(r_i)$$

follows from Theorem 4.1.

## 7 Lecture 7: Full Rank Regression

Model:

$$Y = \beta_0 + \beta_1 x_1 + \cdots + \beta_k x_k + \varepsilon,$$

where  $x_i$  are **predictors**,  $Y$  is the **response**, and  $\varepsilon$  is noise. If we have  $i = 1, \dots, n$  observations, then the model becomes:

$$Y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_k x_{ik} + \varepsilon_i = \mathbf{X}\vec{\beta} + \vec{\varepsilon},$$

where

$$\vec{Y} = \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix}, \quad \mathbf{X} = \begin{pmatrix} 1 & x_{11} & \cdots & x_{1k} \\ 1 & x_{21} & \cdots & x_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n1} & \cdots & x_{nk} \end{pmatrix}, \quad \vec{\beta} = \begin{pmatrix} \beta_0 \\ \vdots \\ \beta_k \end{pmatrix}, \quad \vec{\varepsilon} = \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_n \end{pmatrix}$$

Assumptions:

- (1)  $\mathbb{E}[\varepsilon_i] = 0$ .
- (2)  $\text{Var}(\varepsilon_i) = \sigma^2$ .
- (3)  $\text{Cov}(\varepsilon_i, \varepsilon_j) = 0$  for  $i \neq j$ .

Full rank assumption:

- (1)  $k < n$ ;
- (2)  $\text{rank}(\mathbf{X}) = k + 1$ .

### Method 1: Least Squares Method

$$L = \sum_{i=1}^n \varepsilon_i^2 = \vec{\varepsilon}'\vec{\varepsilon} = (\vec{Y} - \mathbf{X}\vec{\beta})'(\vec{Y} - \mathbf{X}\vec{\beta}).$$

If we minimize with respect to  $\vec{\beta}$ , we get

$$\frac{\partial L}{\partial \vec{\beta}} = \begin{pmatrix} \frac{\partial L}{\partial \beta_0} \\ \vdots \\ \frac{\partial L}{\partial \beta_k} \end{pmatrix} = -2\mathbf{X}\vec{Y} + 2\mathbf{X}'\mathbf{X}\vec{\beta} = 0 \implies \hat{\vec{\beta}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y}.$$

#### THEOREM 7.1

$\hat{\vec{\beta}}$  is an unbiased estimator of  $\vec{\beta}$ .

**Proof:**

$$\begin{aligned} \mathbb{E}[\hat{\vec{\beta}}] &= \mathbb{E}[(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y}] \\ &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}[\vec{Y}] \\ &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}[\mathbf{X}\vec{\beta} + \vec{\varepsilon}] \\ &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\vec{\beta} + \vec{0} \\ &= \vec{\beta}. \end{aligned}$$

**THEOREM 7.2**

If  $\text{Cov}(\vec{Y}, \vec{Y}) = \sigma^2 \mathbf{I}$ , then  $\text{Cov}(\hat{\vec{\beta}}, \hat{\vec{\beta}}) = \sigma^2 (\mathbf{X}'\mathbf{X})^{-1}$ .

**Proof:**

$$\begin{aligned}
 \text{Cov}(\hat{\vec{\beta}}, \hat{\vec{\beta}}) &= \mathbb{E}[(\hat{\vec{\beta}} - \mathbb{E}[\hat{\vec{\beta}}])(\hat{\vec{\beta}} - \mathbb{E}[\hat{\vec{\beta}}])'] \\
 &= \mathbb{E}\left[\left((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y} - (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}[\vec{Y}]\right)\left((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y} - (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}[\vec{Y}]\right)'\right] \\
 &= \mathbb{E}\left[\mathbf{X}'\mathbf{X}^{-1}\mathbf{X}'(\vec{Y} - \mathbb{E}[\vec{Y}])(\vec{Y} - \mathbb{E}[\vec{Y}])'\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\right] \\
 &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}\left[(\vec{Y} - \mathbb{E}[\vec{Y}])(\vec{Y} - \mathbb{E}[\vec{Y}])'\right]\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1} \\
 &= \sigma^2 (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1} \\
 &= \sigma^2 (\mathbf{X}'\mathbf{X})^{-1}.
 \end{aligned}$$

**Estimation of  $\sigma^2$** 

- Residual:

$$\begin{aligned}
 (\vec{Y} - \hat{\vec{Y}}) &= \hat{\vec{\epsilon}} \\
 &= (\vec{Y} - \mathbf{X}\hat{\vec{\beta}}) \\
 &= (\vec{Y} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y}) \\
 &= (\mathbf{I} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}')\vec{Y}.
 \end{aligned}$$

- Let  $\mathbf{H} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$  be the **hat matrix**.
- Since  $\mathbf{H}$  is idempotent, we may write

$$\begin{aligned}
 \text{SSE} &= \hat{\vec{\epsilon}}'\hat{\vec{\epsilon}} \\
 &= \|\vec{Y} - \hat{\vec{Y}}\|^2 \\
 &= (\vec{Y} - \hat{\vec{Y}})'(\vec{Y} - \hat{\vec{Y}}) \\
 &= \vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}.
 \end{aligned}$$

**THEOREM 7.3**

$$S^2 = \frac{\text{SSE}}{n - (k + 1)}$$

is an unbiased estimator of  $\sigma^2$ .

**Proof:**

$$\begin{aligned}
\mathbb{E}[S^2] &= \frac{1}{n - (k + 1)} \mathbb{E}[\vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}] \\
&= \frac{1}{n - (k + 1)} [\text{tr}((\mathbf{I} - \mathbf{H})\boldsymbol{\Sigma}) + \vec{\mu}'(\mathbf{I} - \mathbf{H})\vec{\mu}] \\
&= \frac{\sigma^2}{n - (k + 1)} \text{tr}(\mathbf{I} - \mathbf{H}) \\
&= \frac{\sigma^2}{n - (k + 1)} (\text{tr}(\mathbf{I}) - \text{tr}(\mathbf{H})) \\
&= \frac{\sigma^2}{n - (k + 1)} (\text{tr}(\mathbf{I}) - \text{rank}(\mathbf{H})) \\
&= \frac{\sigma^2}{n - (k + 1)} (n - (k + 1)) \\
&= \sigma^2.
\end{aligned}$$

Note that  $\vec{\mu} = \mathbf{X}\vec{\beta} \implies \vec{\mu}' = \vec{\beta}'\mathbf{X}'$ , so

$$\begin{aligned}
\mathbf{X}'(\mathbf{I} - \mathbf{H})\mathbf{X} &= \mathbf{X}'\mathbf{X} - \mathbf{X}'\mathbf{H}\mathbf{X} \\
&= \mathbf{X}'\mathbf{X} - \mathbf{X}'\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X} \\
&= \mathbf{O}.
\end{aligned}$$

## Maximum Likelihood Estimators for $\vec{\beta}$ and $\sigma^2$

### THEOREM 7.4

If  $\vec{Y} \sim MN(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I}_n)$ , where  $\mathbf{X}$  is  $n \times (k + 1)$  of rank  $k + 1 < n$ , then the maximum likelihood estimators of  $\vec{\beta}$  and  $\sigma^2$  are

$$\hat{\vec{\beta}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y}, \quad \hat{\sigma}^2 = \frac{SSE}{n} = \frac{(\vec{Y} - \mathbf{X}\hat{\vec{\beta}})'(\vec{Y} - \mathbf{X}\hat{\vec{\beta}})}{n}.$$

**Proof:** The likelihood function is given by the multivariate normal density

$$\begin{aligned}
L(\vec{\beta}, \sigma^2) &= f(\vec{Y}; \vec{\beta}, \sigma^2) \\
&= \frac{1}{(2\pi)^{n/2} |\sigma^2\mathbf{I}|^{1/2}} \exp\left\{-\frac{(\vec{Y} - \mathbf{X}\vec{\beta})'(\sigma^2\mathbf{I})^{-1}(\vec{Y} - \mathbf{X}\vec{\beta})}{2}\right\} \\
&= (2\pi\sigma^2)^{-n/2} \exp\left\{-\frac{(\vec{Y} - \mathbf{X}\vec{\beta})'(\vec{Y} - \mathbf{X}\vec{\beta})}{2\sigma^2}\right\}.
\end{aligned}$$

The log-likelihood function is

$$\begin{aligned}
\ell(\vec{\beta}, \sigma^2) &= \ln(L(\vec{\beta}, \sigma^2)) \\
&= -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} (\vec{Y} - \mathbf{X}\vec{\beta})'(\vec{Y} - \mathbf{X}\vec{\beta}) \\
&= -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} (\vec{Y}'\vec{Y} - 2\vec{Y}'\mathbf{X}\vec{\beta} + \vec{\beta}'\mathbf{X}'\mathbf{X}\vec{\beta}).
\end{aligned}$$

Taking the derivative with respect to  $\vec{\beta}$  yields

$$\begin{aligned} \vec{0} &= \frac{\partial \ell(\vec{\beta}, \sigma^2)}{\partial \vec{\beta}} = -\frac{1}{2\sigma^2}(-2\mathbf{X}'\vec{Y} + 2\mathbf{X}'\mathbf{X}\vec{\beta}) \\ \vec{0} &= 2\mathbf{X}'\vec{Y} - 2\mathbf{X}'\mathbf{X}\vec{\beta} \\ \mathbf{X}'\mathbf{X}\vec{\beta} &= \mathbf{X}'\vec{Y} \\ \hat{\vec{\beta}} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\vec{Y}. \end{aligned}$$

Taking the derivative with respect to  $\sigma^2$  yields

$$\begin{aligned} 0 &= \frac{\partial \ell(\vec{\beta}, \sigma^2)}{\partial \sigma^2} = -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4}(\vec{Y} - \mathbf{X}\vec{\beta})'(\vec{Y} - \mathbf{X}\vec{\beta}) \\ \frac{n(2\sigma^4)}{2\sigma^2} &= (\vec{Y} - \mathbf{X}\vec{\beta})'(\vec{Y} - \mathbf{X}\vec{\beta}) \\ \hat{\sigma}^2 &= \frac{(\vec{Y} - \mathbf{X}\hat{\vec{\beta}})'(\vec{Y} - \mathbf{X}\hat{\vec{\beta}})}{n}. \end{aligned}$$

## Properties of $\hat{\vec{\beta}}$ and $\hat{\sigma}^2$

### THEOREM 7.5

If  $\vec{Y} \sim MN(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I}_n)$ , where  $\mathbf{X}$  is  $n \times (k+1)$  of rank  $k+1 < n$ , and  $\vec{\beta} = (\beta_0, \beta_1, \dots, \beta_k)'$ , then the maximum likelihood estimators of  $\vec{\beta}$  and  $\hat{\sigma}^2$  given in Theorem 7.1 have the following distributional properties:

- (1)  $\hat{\vec{\beta}} \sim MN(\vec{\beta}, \sigma^2(\mathbf{X}'\mathbf{X})^{-1})$ .
- (2)  $\frac{n\hat{\sigma}^2}{\sigma^2} \sim \chi^2(n - (k+1))$ .
- (3)  $\hat{\vec{\beta}}$  and  $\hat{\sigma}^2$  are independent.

**Proof:**

- (1) Note that  $\vec{Y} \sim MN(\vec{\mu}, \Sigma) \implies \mathbf{A}\vec{Y} \sim MN(\mathbf{A}\vec{\mu}, \mathbf{A}\Sigma\mathbf{A}')$ . Let  $\mathbf{A} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ ,  $\vec{\mu} = \mathbf{X}\vec{\beta}$ , and  $\Sigma = \sigma^2\mathbf{I}_n$ . Now,

$$\begin{aligned} \mathbf{A}\vec{Y} = \hat{\vec{\beta}} &\sim MN\left((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}\vec{\beta}, (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\sigma^2\mathbf{I}_n\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\right) \\ &\sim MN(\vec{\beta}, \sigma^2(\mathbf{X}'\mathbf{X})^{-1}). \end{aligned}$$

- (2) Note that

$$\begin{aligned} \frac{n\hat{\sigma}^2}{\sigma^2} &= \frac{\text{SSE}}{\sigma^2} \\ &= \frac{\vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}}{\sigma^2} \\ &= \vec{W}'(\mathbf{I} - \mathbf{H})\vec{W}, \end{aligned}$$

where  $\vec{W} = \frac{\vec{Y}}{\sigma} \sim MN\left(\frac{\mathbf{X}\vec{\beta}}{\sigma}, \mathbf{I}\right)$ . It follows from Theorem 4.1 that

$$\frac{n\hat{\sigma}^2}{\sigma^2} \sim \chi^2(r, \lambda),$$

with  $r = \text{rank}(\mathbf{I} - \mathbf{H}) = \text{tr}(\mathbf{I} - \mathbf{H}) = n - (k + 1)$  and

$$\begin{aligned}\lambda &= \frac{1}{2} \vec{\mu}' \mathbf{A} \vec{\mu} \\ &= \frac{1}{2} \left( \frac{\mathbf{X} \vec{\beta}}{\sigma} \right)' (\mathbf{I} - \mathbf{H}) \frac{\mathbf{X} \vec{\beta}}{\sigma} \\ &= \frac{1}{2\sigma^2} \vec{\beta}' [\mathbf{X}'(\mathbf{I} - \mathbf{H})\mathbf{X}] \vec{\beta} \\ &= 0,\end{aligned}$$

where  $\vec{\mu} = \mathbb{E}[\vec{W}]$  and  $\mathbf{A} = \mathbf{I} - \mathbf{H}$ .

(3) Note that  $\vec{\beta} = (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\vec{Y}$  and

$$\hat{\sigma}^2 = \frac{\text{SSE}}{n} = \frac{1}{n} \vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y} = \vec{Y}' \left( \frac{\mathbf{I} - \mathbf{H}}{n} \right) \vec{Y}.$$

Let  $\vec{Y} = \vec{X}$ ,  $\mathbf{B} = (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}$ , and  $\mathbf{A} = \frac{\mathbf{I} - \mathbf{H}}{n}$ . Relabelling,

$$\begin{aligned}\vec{\beta} &= \mathbf{B}\vec{X}, \\ \hat{\sigma}^2 &= \vec{X}' \mathbf{A} \vec{X}.\end{aligned}$$

Now,

$$\begin{aligned}(\mathbf{X}'\mathbf{X})^{-1} \mathbf{X} \left( \frac{\mathbf{I} - \mathbf{H}}{n} \right) &= \frac{1}{n} [(\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}' - (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'] \\ &= 0.\end{aligned}$$

The result follows from Theorem 5.1.

## 8 Lecture 8: Test of Overall Regression

**DEFINITION 8.1: Sum of Squares Total, Residual, Error**

$$\begin{aligned}\bar{Y} &= \frac{1}{n} \sum_{i=1}^n Y_i, \\ \text{SST} &= \sum_{i=1}^n (Y_i - \bar{Y})^2, \\ \text{SSR} &= \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2, \\ \text{SSE} &= \sum_{i=1}^n (Y_i - \hat{Y}_i)^2.\end{aligned}$$

SST is the **sum of squares total**, SSR is the **sum of squares residual**, and SSE is the **sum of squares error**.



**THEOREM 8.1**

$$(i) \text{ SST} = \vec{Y}'(\mathbf{I} - \frac{1}{n}\mathbf{J})\vec{Y}.$$

$$(ii) \text{ SSR} = \vec{Y}'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{Y}.$$

$$(iii) \text{ SSE} = \vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}.$$

Hence,  $\text{SST} = \text{SSR} + \text{SSE}$ .

**Proof:**

(i) Sum of Squares Total:

$$\begin{aligned} \text{SST} &= \sum_{i=1}^n (Y_i - \bar{Y})^2 \\ &= (\vec{Y} - \bar{Y}\vec{j})'(\vec{Y} - \bar{Y}\vec{j}) \\ &= \vec{Y}'\vec{Y} - 2\bar{Y}\vec{Y}'\vec{j} + \bar{Y}^2\vec{j}'\vec{j} \\ &= \vec{Y}'\vec{Y} - 2n\bar{Y}^2 + n\bar{Y}^2 \\ &= \vec{Y}'\vec{Y} - n\bar{Y}^2 \\ &= \vec{Y}'\vec{Y} - \frac{1}{n}\vec{Y}'\vec{j}\vec{j}'\vec{Y} \\ &= \vec{Y}'(\mathbf{I} - \frac{1}{n}\mathbf{J})\vec{Y}. \end{aligned}$$

(ii) Sum of Squares Regression:

$$\begin{aligned} \text{SSR} &= \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2 \\ &= (\hat{\vec{Y}} - \bar{Y}\vec{j})'(\hat{\vec{Y}} - \bar{Y}\vec{j}) \\ &= (\mathbf{H}\vec{Y} - \bar{Y}\vec{j})'(\mathbf{H}\vec{Y} - \bar{Y}\vec{j}) \\ &= \vec{Y}'\mathbf{H}\mathbf{H}\vec{Y} - 2\bar{Y}\vec{Y}'\mathbf{H}\vec{j} + \bar{Y}^2\vec{j}'\vec{j} \\ &= \vec{Y}'\mathbf{H}^2\vec{Y} - \frac{2}{n}\vec{Y}'\mathbf{H}\vec{j}\vec{j}'\vec{Y} + \frac{1}{n}\vec{Y}'\mathbf{J}\vec{Y} \\ &= \vec{Y}'\mathbf{H}\vec{Y} - \frac{2}{n}\vec{Y}'\mathbf{H}\mathbf{J}\vec{Y} + \frac{1}{n}\vec{Y}'\mathbf{J}\vec{Y} \\ &= \vec{Y}'\mathbf{H}\vec{Y} - \frac{2}{n}\vec{Y}'\mathbf{J}\vec{Y} + \frac{1}{n}\vec{Y}'\mathbf{J}\vec{Y} \\ &= \vec{Y}'\mathbf{H}\vec{Y} - \frac{1}{n}\vec{Y}'\mathbf{J}\vec{Y} \\ &= \vec{Y}'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{Y} \end{aligned}$$

since  $\mathbf{H}^2 = \mathbf{H}$ ,  $\mathbf{H}\mathbf{X} = \mathbf{X}$ , and  $\mathbf{H}\vec{j} = \vec{j} \implies \mathbf{H}\mathbf{J} = \mathbf{J}$ .

(iii) Sum of Squares Error:

$$\begin{aligned} \text{SSE} &= \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 \\ &= (\vec{Y} - \hat{\vec{Y}})'(\vec{Y} - \hat{\vec{Y}}) \\ &= (\vec{Y} - \mathbf{H}\vec{Y})'(\vec{Y} - \mathbf{H}\vec{Y}) \\ &= ((\mathbf{I} - \mathbf{H})\vec{Y})'(\mathbf{I} - \mathbf{H})\vec{Y} \\ &= \vec{Y}'(\mathbf{I} - \mathbf{H})^2\vec{Y} \\ &= \vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y} \end{aligned}$$

since  $(\mathbf{I} - \mathbf{H})^2 = \mathbf{I} - \mathbf{H}$ .

Therefore,

$$\mathbf{I} - \mathbf{H} + \mathbf{H} - \frac{1}{n}\mathbf{J} = \mathbf{I} - \frac{1}{n}\mathbf{J} \implies \text{SST} = \text{SSR} + \text{SSE}.$$

### THEOREM 8.2

(1)  $\text{SSR}/\sigma^2 \sim \chi^2(k, \lambda)$  with

$$\begin{aligned} \lambda &= \frac{1}{2\sigma^2} (\mathbf{X}\vec{\beta})' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \mathbf{X}\vec{\beta} \\ &= \frac{1}{2\sigma^2} (\mathbf{X}_1\vec{\beta}_1)' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \mathbf{X}_1\vec{\beta}_1, \end{aligned}$$

where

$$\vec{\beta}_1 = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_k \end{pmatrix}, \quad \mathbf{X}_1 = \begin{pmatrix} X_{11} & \cdots & X_{1k} \\ \vdots & \ddots & \vdots \\ X_{n1} & \cdots & X_{nk} \end{pmatrix}.$$

(2)  $\text{SSR}$  and  $\text{SSE}$  are independent.

**Proof:**

(1) First result:

$$\frac{\text{SSR}}{\sigma^2} = \frac{\vec{Y}'}{\sigma} (\mathbf{H} - \frac{1}{n}\mathbf{J}) \frac{\vec{Y}}{\sigma}$$

with

$$\frac{\vec{Y}}{\sigma} \sim \text{MN}\left(\frac{\mathbf{X}\vec{\beta}}{\sigma}, \mathbf{I}\right).$$

Since

$$\begin{aligned} (\mathbf{H} - \frac{1}{n}\mathbf{J})^2 &= \mathbf{H}^2 - 2\frac{1}{n}\mathbf{H}\mathbf{J} + \frac{1}{n^2}\mathbf{J}^2 \\ &= \mathbf{H} - 2\frac{1}{n}\mathbf{J} + \frac{1}{n}\mathbf{J} \\ &= \mathbf{H} - \frac{1}{n}\mathbf{J} \end{aligned}$$

it follows from Theorem 4.1 that  $\text{SSR}/\sigma^2 \sim \chi^2(r, \lambda)$  with

$$r = \text{rank}(\mathbf{H} - \frac{1}{n}\mathbf{J}) = \text{tr}(\mathbf{H} - \frac{1}{n}\mathbf{J}) = k + 1 - 1 = k$$

and

$$\lambda = \frac{1}{n} \left( \frac{\mathbf{X}\vec{\beta}}{\sigma} \right)' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \frac{\mathbf{X}\vec{\beta}}{\sigma}.$$

Write

$$\vec{\beta} = \begin{pmatrix} \beta_0 \\ \vec{\beta}_1 \end{pmatrix}, \quad \mathbf{X} = \begin{pmatrix} \vec{j} & \mathbf{X}_1 \end{pmatrix}.$$

Then,  $\mathbf{X}\vec{\beta} = \beta_0\vec{j} + \mathbf{X}_1\vec{\beta}_1$  implies

$$\begin{aligned} \lambda &= \frac{1}{2} \left( \frac{\beta_0\vec{j} + \mathbf{X}_1\vec{\beta}_1}{\sigma} \right)' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \frac{\beta_0\vec{j} + \mathbf{X}_1\vec{\beta}_1}{\sigma} \\ &= \frac{1}{2\sigma^2} \left[ (\mathbf{X}_1\vec{\beta}_1)' (\mathbf{H} - \frac{1}{n}\mathbf{J}) (\mathbf{X}_1\vec{\beta}_1) + \beta_0^2 \vec{j}' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \vec{j} + 2\beta_0 (\mathbf{H} - \frac{1}{n}\mathbf{J}) \vec{j} \right] \\ &= \frac{1}{2\sigma^2} (\mathbf{X}_1\vec{\beta}_1)' (\mathbf{H} - \frac{1}{n}\mathbf{J}) \mathbf{X}_1\vec{\beta}_1 \end{aligned}$$

since

$$(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{j} = \mathbf{H}\vec{j} - \frac{1}{n}\mathbf{J}\vec{j} = \vec{j} - \frac{1}{n}n\vec{j} = \vec{0}.$$

(2) Note that

$$\text{SSR} = \vec{Y}'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{Y} = \vec{Y}'\mathbf{A}\vec{Y}.$$

$$\text{SSE} = \vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y} = \vec{Y}'\mathbf{B}\vec{Y}.$$

Note that  $\vec{Y} \sim \text{MN}(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I})$ . Note that  $\mathbf{H} - \frac{1}{n}\mathbf{J}$  and  $\mathbf{I} - \mathbf{H}$  are symmetric matrices of the same dimension.

$$\begin{aligned} \mathbf{A}\sigma^2\mathbf{I}\mathbf{B} &= \sigma^2\mathbf{A}\mathbf{B} \\ &= \sigma^2(\mathbf{H} - \frac{1}{n}\mathbf{J})(\mathbf{I} - \mathbf{H}) \\ &= \sigma^2(\mathbf{H} - \frac{1}{n}\mathbf{J} - \mathbf{H}^2 + \frac{1}{n}\mathbf{J}\mathbf{H}) \\ &= \sigma^2(\mathbf{H} - \frac{1}{n}\mathbf{J} - \mathbf{H} + \frac{1}{n}\mathbf{J}) \\ &= \sigma^2\mathbf{O} \\ &= \mathbf{O}. \end{aligned}$$

Therefore, SSR and SSE are independent by Theorem 5.2.

#### REMARK

By Theorem 7.4, we have  $\text{SSE}/\sigma^2 \sim \chi^2(n - (k + 1))$ , and  $\text{SST}/\sigma^2 \sim \chi^2(n - 1, \lambda)$ .

#### REMARK

$$\begin{aligned} \mathbb{E}[\text{SSR}] &= \sigma^2 \mathbb{E}\left[\frac{\text{SSR}}{\sigma^2}\right] \\ &= \sigma^2 \sum_{i=1}^k \mathbb{E}[X_i^2] \quad \text{where } X_i \sim \mathcal{N}(\mu_i, 1) \\ &= \sigma^2 \left( \sum_{i=1}^n (\text{Var}(X_i) + \mu_i^2) \right) \\ &= \sigma^2(k + 2\lambda) \\ &= k\sigma^2 + 2\sigma^2\lambda \\ &= k\sigma^2 + (\mathbf{X}_1\vec{\beta}_1)'(\mathbf{H} - \frac{1}{n}\mathbf{J})\mathbf{X}_1\vec{\beta}_1. \end{aligned}$$

### ANOVA Table for Hypothesis Test of $\mathbf{H}_0: \vec{\beta}_1 = \vec{0}$ versus $\mathbf{H}_A: \vec{\beta}_1 \neq \vec{0}$

| Source of Variation    | Degrees of Freedom | Sum of Squares | Mean Square | Expected Mean Square  |
|------------------------|--------------------|----------------|-------------|---|
| Due to $\vec{\beta}_1$ | $k$                | SSR            | MSR         | $\sigma^2 + \frac{1}{k}(\mathbf{X}_1\vec{\beta}_1)'(\mathbf{H} - \frac{1}{n}\mathbf{J})\mathbf{X}_1\vec{\beta}_1$ |
| Error                  | $n - (k + 1)$      | SSE            | MSE         | $\sigma^2$  |
| Total                  | $n - 1$            | SST            |             |   |

Note that under  $H_0$ ,  $\frac{1}{k}(\mathbf{X}_1 \vec{\beta}_1)'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{X}_1 \vec{\beta}_1 = 0$ .

$$\begin{aligned}\mathbb{E}[\text{MSR}] &= \mathbb{E}\left[\frac{\text{SSR}}{k}\right] \\ &= \frac{1}{k}\left[\sigma^2 k + (\mathbf{X}_1 \vec{\beta}_1)'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{X}_1 \vec{\beta}_1\right] \\ &= \sigma^2 + \frac{1}{k}(\mathbf{X}_1 \vec{\beta}_1)'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{X}_1 \vec{\beta}_1. \\ \mathbb{E}[\text{MSE}] &= \mathbb{E}\left[\frac{\text{SSE}}{n - (k + 1)}\right] \\ &= \frac{(n - (k + 1))\sigma^2}{n - (k + 1)} \\ &= \sigma^2.\end{aligned}$$

Test Statistic:

$$F = \frac{\text{SSR}/k}{\text{SSE}/(n - (k + 1))} \sim F(k, n - (k + 1)).$$

Reject  $H_0$  if  $F > F_\alpha(k, n - (k + 1))$ . If  $H_0$  holds, then  $\lambda = 0$  implies  $\text{SSR} \sim \chi^2(k)$  and  $\text{SSE} \sim \chi^2(n - (k + 1))$ . Furthermore, note that if  $X$  and  $Y$  are independent, then  $f(X)$  and  $g(Y)$  are independent, so

$$\begin{aligned}\mathbb{E}[F] &= \mathbb{E}\left[\frac{\text{SSR}/k}{\text{SSE}/(n - (k + 1))}\right] \\ &= \frac{n - (k + 1)}{k} \mathbb{E}[\text{SSR}] \mathbb{E}\left[\frac{1}{\text{SSE}}\right]. \\ \mathbb{E}\left[\frac{\text{SSR}}{k}\right] &= \sigma^2 \iff \mathbb{E}[\text{SSR}] = k\sigma^2.\end{aligned}$$

Hold tight for the hard part (you can skip this part if you know the mean of the *inverse-chi-squared distribution*),

$$\begin{aligned}\mathbb{E}\left[\frac{\sigma^2}{\text{SSE}}\right] &= \int_0^\infty \frac{1}{y} \frac{1}{2^{(n-k-1)/2} \Gamma\left(\frac{n-k-1}{2}\right)} y^{(n-k-1)/2-1} e^{-y/2} dy \\ &= 2^{-(n-k-1)/2} \frac{\Gamma\left(\frac{n-k-1}{2} - 1\right)}{\Gamma\left(\frac{n-k-1}{2}\right)} \times \\ &\quad \underbrace{2^{(n-k-1)/2-1} \frac{1}{\Gamma\left(\frac{n-k-1}{2} - 1\right) 2^{(n-k-1)/2-1}} \int_0^\infty y^{((n-k-1)/2-1)-1} e^{-y/2} dy}_{=1 \text{ by Gamma distribution}} \\ &= \frac{1}{2} \frac{1}{(n-k-1)/2-1} \\ &= \frac{1}{n-k-3}.\end{aligned}$$

Hence,

$$\mathbb{E}\left[\frac{1}{\text{SSE}}\right] = \frac{1}{\sigma^2(n-k-3)}.$$

Therefore,

$$\mathbb{E}[F] = \frac{n-k-1}{k} \mathbb{E}[\text{SSR}] \mathbb{E}\left[\frac{1}{\text{SSE}}\right] = \frac{n-k-1}{k} k\sigma^2 \frac{1}{\sigma^2(n-k-3)} = \frac{n-k-1}{n-k-3}.$$

## 9 Lecture 9: Lack of Fit

Consider the case of studying blood pressure and its relationship to height and weight. Clearly, people of the same height and weight can have different blood pressures. In other words, the same predictor values may correspond to different response values. This type of variation is called pure error. To detect poor model fit, we would need to distinguish between variation caused by the model and pure error.

### General Framework

Let  $m \geq 1$  and  $n_1, \dots, n_m \geq 1$  such that  $\sum_{i=1}^m n_i = n$ . For  $i = 1, \dots, m$ , we have

$$Y_{ir} = \beta_0 + \beta_1 x_{1i} + \dots + \beta_k x_{ik} + \varepsilon_{ir}, \quad r = 1, \dots, n_i.$$

In matrix notation, we write  $\mathbf{Y} = \mathbf{X}\vec{\beta}$ , where  $\vec{\beta} = \begin{pmatrix} \beta_0 \\ \vdots \\ \beta_k \end{pmatrix}$  in the usual way, and

$$\begin{aligned} \vec{Y}' &= (Y_{11} \quad \dots \quad Y_{1n_1} \quad \dots \quad Y_{m1} \quad \dots \quad Y_{mn_m}), \\ \mathbf{X} &= \begin{pmatrix} n_1 \begin{pmatrix} 1 & x_{11} & \dots & x_{1k} \\ \vdots & \ddots & \ddots & \vdots \\ 1 & x_{11} & \dots & x_{1k} \end{pmatrix} \\ \vdots \\ n_m \begin{pmatrix} 1 & x_{m1} & \dots & x_{mk} \\ \vdots & \ddots & \ddots & \vdots \\ 1 & x_{m1} & \dots & x_{mk} \end{pmatrix} \end{pmatrix}, \\ \vec{\varepsilon}' &= (\varepsilon_{11} \quad \dots \quad \varepsilon_{1n_1} \quad \dots \quad \varepsilon_{m1} \quad \dots \quad \varepsilon_{mn_m}). \end{aligned}$$

We write  $Y_{ij}$  for  $i = 1, \dots, m$  ( $m$  groups) and  $j = 1, \dots, n_i$  (number of observations in group  $i$ ). The sample average of group  $i$  is defined by

$$\bar{Y}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}, \quad i = 1, \dots, m.$$

The fitted values are  $\hat{Y} = \mathbf{X}\hat{\beta}$ , so  $\hat{Y}_{ij}$  is the same for all  $j = 1, \dots, n_i$ , hence we may write  $\hat{Y}_{ij}$  as  $\hat{Y}_i$ .

$$\begin{aligned} \text{SSE} &= \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \hat{Y}_{ij})^2 \\ &= \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i + \bar{Y}_i - \hat{Y}_{ij})^2 \\ &= \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2 + \sum_{i=1}^m \sum_{j=1}^{n_i} (\hat{Y}_{ij} - \bar{Y}_i)^2 - 2 \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)(\bar{Y}_i - \hat{Y}_{ij}) \\ &= \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2 + \sum_{i=1}^m \sum_{j=1}^{n_i} (\hat{Y}_{ij} - \bar{Y}_i)^2 \\ &= \sum_{i=1}^m \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2 + \sum_{i=1}^m n_i (\hat{Y}_i - \bar{Y}_i)^2 \\ &= \text{SSPE} + \text{SSLF} \end{aligned}$$

since  $\hat{Y}_{ij}$  is independent of  $j$ . Therefore,

$$\text{SST} = \text{SSR} + \text{SSE} = \text{SSR} + \text{SSPE} + \text{SSLF}.$$

- Degrees of freedom of SSLF:  $m - (k + 1)$ .
- Degrees of freedom of SSPE:  $(n_1 - 1) + \dots + (n_m - 1) = n - m$ .

The first test is a test of linear relationship, but if we wanted to determine how good that relationship is, we will need the following hypothesis test. If the linear model fits well, then SSLF should be small.

- $H_0$ : The model is adequate.
- $H_A$ : The model is not adequate.

Test statistic:

$$F = \frac{\text{SSLF}/(m - k - 1)}{\text{SSPE}/(n - m)} \sim F(m - k - 1, n - m).$$

If we reject  $H_0$ , that means there's too much variation within the group. Reject  $H_0$  when  $F > F_\alpha(m - k - 1, n - m)$ .

| Source of Variation    | Degrees of Freedom | Sum of Squares | Mean Square | $F$       |
|------------------------|--------------------|----------------|-------------|-----------|
| Due to $\vec{\beta}_1$ | $k$                | SSR            | MSR         | MSR/MSE   |
| Error                  | $n - (k + 1)$      | SSE            | MSE         |           |
| Lack of Fit            | $m - k - 1$        | SSLF           | MSLF        | MSLF/MSPE |
| Pure Error             | $n - m$            | SSPE           | MSPE        |           |
| Total                  | $n - 1$            | SST            |             |           |

## Selection of Predictors

We observe that the number of predictors always improves the estimates, but becomes less efficient. To find a reasonable number of predictors, one needs to compare models by adding or dropping predictors.

- Partition  $\vec{\beta}$  as  $\vec{\beta} = \begin{pmatrix} \beta_0 \\ \vdots \\ \beta_k \end{pmatrix} = \begin{pmatrix} \vec{\beta}_I \\ \vec{\beta}_{II} \end{pmatrix}$ , where  $\vec{\beta}_I = \begin{pmatrix} \beta_0 \\ \vdots \\ \beta_\ell \end{pmatrix}$  and  $\vec{\beta}_{II} = \begin{pmatrix} \beta_{\ell+1} \\ \vdots \\ \beta_k \end{pmatrix}$  for  $1 \leq \ell < k$ .
- Partition  $\mathbf{X}$  as  $\mathbf{X} = (\mathbf{X}_I \quad \mathbf{X}_{II})$ , where  $\mathbf{X}_I \in \mathbb{R}^{n \times (\ell+1)}$  and  $\mathbf{X}_{II} \in \mathbb{R}^{n \times (k-\ell)}$  for  $1 \leq \ell < k$ .

The **full model** is

$$\vec{Y} = \mathbf{X}\vec{\beta} + \vec{\varepsilon} = (\mathbf{X}_I \quad \mathbf{X}_{II}) \begin{pmatrix} \vec{\beta}_I \\ \vec{\beta}_{II} \end{pmatrix} = \mathbf{X}_I\vec{\beta}_I + \mathbf{X}_{II}\vec{\beta}_{II}.$$

The **reduced model** is

$$\vec{Y} = \mathbf{X}_I\vec{\beta}_I + \vec{\varepsilon}^*.$$

Let

- $\mathbf{H} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ .
- $\mathbf{H}_I = \mathbf{X}_I(\mathbf{X}'_I\mathbf{X}_I)^{-1}\mathbf{X}'_I$ .

Define

- $\text{SS}(\vec{\beta}) = \text{SSR}(\text{full}) = \vec{Y}'(\mathbf{H} - \frac{1}{n}\mathbf{J})\vec{Y}$ .
- $\text{SS}(\vec{\beta}_I) = \text{SSR}(\text{reduced}) = \vec{Y}'(\mathbf{H}_I - \frac{1}{n}\mathbf{J})\vec{Y}$ .
- $\text{SS}(\vec{\beta}_{II} | \vec{\beta}_I) = \text{SS}(\vec{\beta}) - \text{SS}(\vec{\beta}_I) = \vec{Y}'(\mathbf{H} - \mathbf{H}_I)\vec{Y}$ .

Comparing the full model and the reduced model, we test  $H_0: \vec{\beta}_{II} = \vec{0}$  versus  $H_A: \vec{\beta}_{II} \neq \vec{0}$ . Under  $H_0$ ,  $x_{\ell+1}, \dots, x_k$  do not add predictive value to the model that includes  $x_1, \dots, x_\ell$  already.

**THEOREM 9.1** $\mathbf{H} - \mathbf{H}_1$  is idempotent.**Proof:** Assignment 2.

Model misspecification:

- Leaving out  $\vec{\beta}_{II}$  when it should be included, results in underfitting.
- Including  $\vec{\beta}_{II}$  when it should be dropped, results in overfitting.

LECTURE 10  
9th February**10 Lecture 10: Determination of Predictors and Generalized Inverse****THEOREM 10.1**Let  $\vec{Y} \sim MN(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I})$ ,  $\mathbf{H} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ ,  $\mathbf{H}_1 = \mathbf{X}_I(\mathbf{X}'_I\mathbf{X}_I)^{-1}\mathbf{X}'_I$ . Then,

(1)  $\vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}/\sigma^2 = SSE/\sigma^2 \sim \chi^2(n - k - 1)$ .

(2)  $\vec{Y}'(\mathbf{H} - \mathbf{H}_1)\vec{Y}/\sigma^2 = \frac{SS(\vec{\beta}_{II}|\vec{\beta}_I)}{\sigma^2} \sim \chi^2(k - \ell, \tilde{\lambda})$ , where

$$\tilde{\lambda} = \vec{\beta}'_{II}\mathbf{X}'_{II}(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II}/2\sigma^2.$$

(3)  $\vec{Y}'(\mathbf{I} - \mathbf{H})\vec{Y}$  and  $\vec{Y}'(\mathbf{H} - \mathbf{H}_1)\vec{Y}$  are independent.

**Proof:**

(1) Earlier proof.

(2)

$$\begin{aligned} \vec{Y}'(\mathbf{H} - \mathbf{H}_1)\vec{Y}/\sigma^2 &= \frac{SS(\text{full} | \text{reduced})}{\sigma^2} \\ &= \frac{SS(\vec{\beta}_{II} | \vec{\beta}_I)}{\sigma^2}. \end{aligned}$$

By Theorem 9.1,  $\mathbf{H} - \mathbf{H}_1$  is idempotent, so

$$\vec{Y}'(\mathbf{H} - \mathbf{H}_1)\vec{Y} \sim \chi^2(r, \tilde{\lambda})$$

 $r = \text{rank}(\mathbf{H} - \mathbf{H}_1)$ , and

$$\tilde{\lambda} = \frac{1}{2\sigma^2}(\mathbf{X}\vec{\beta})'(\mathbf{H} - \mathbf{H}_1)\mathbf{X}\vec{\beta}.$$

By direct calculation, we have

$$\begin{aligned} r &= \text{rank}(\mathbf{H} - \mathbf{H}_1) \\ &= \text{tr}(\mathbf{H} - \mathbf{H}_1) && \text{since } \mathbf{H} - \mathbf{H}_1 \text{ is idempotent} \\ &= \text{tr}(\mathbf{H}) - \text{tr}(\mathbf{H}_1) \\ &= \text{tr}(\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}') - \text{tr}(\mathbf{X}_I(\mathbf{X}'_I\mathbf{X}_I)^{-1}\mathbf{X}'_I) \\ &= \text{tr}(\mathbf{I}_{k+1}) - \text{tr}(\mathbf{I}_{\ell+1}) && \text{cyclic property} \\ &= k - \ell. \end{aligned}$$

Noting that  $\mathbf{H}\mathbf{X} = \mathbf{X}$ ,  $\mathbf{H}_1\mathbf{X}_I = \mathbf{X}_I$ , and

$$\mathbf{X}\vec{\beta} = \mathbf{X}_I\vec{\beta}_I + \mathbf{X}_{II}\vec{\beta}_{II}$$

it follows that

$$\begin{aligned} 2\sigma^2\tilde{\lambda} &= (\mathbf{X}\vec{\beta})'(\mathbf{H} - \mathbf{H}_1)\mathbf{X}\vec{\beta} \\ &= (\mathbf{X}\vec{\beta})'\mathbf{H}\mathbf{X}\vec{\beta} - (\mathbf{X}\vec{\beta})'\mathbf{H}_1\mathbf{X}\vec{\beta} \\ &= (\mathbf{X}\vec{\beta})'(\mathbf{X}\vec{\beta}) - (\mathbf{X}\vec{\beta})'(\mathbf{H}_1\mathbf{X}_I\vec{\beta}_I + \mathbf{H}_1\mathbf{X}_{II}\vec{\beta}_{II}) \\ &= (\mathbf{X}\vec{\beta})'[\mathbf{X}\vec{\beta} - \mathbf{H}_1\mathbf{X}_I\vec{\beta}_I - \mathbf{H}_1\mathbf{X}_{II}\vec{\beta}_{II}] \\ &= (\mathbf{X}\vec{\beta})'[\mathbf{X}_I\vec{\beta}_I + \mathbf{X}_{II}\vec{\beta}_{II} - \mathbf{X}_I\vec{\beta}_I - \mathbf{H}_1\mathbf{X}_{II}\vec{\beta}_{II}] \\ &= (\mathbf{X}\vec{\beta})'(\mathbf{X}_{II}\vec{\beta}_{II} - \mathbf{H}_1\mathbf{X}_{II}\vec{\beta}_{II}) \\ &= (\mathbf{X}\vec{\beta})'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II} \\ &= (\mathbf{X}_I\vec{\beta}_I + \mathbf{X}_{II}\vec{\beta}_{II})'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II} \\ &= (\mathbf{X}_{II}\vec{\beta}_{II})'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II} + (\mathbf{X}_I\vec{\beta}_I)'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II}. \end{aligned}$$

It remains to show that  $(\mathbf{X}_I\vec{\beta}_I)'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II} = \mathbf{O}$ .

$$\begin{aligned} (\mathbf{X}_I\vec{\beta}_I)'(\mathbf{I} - \mathbf{H}_1)\mathbf{X}_{II}\vec{\beta}_{II} &= (\mathbf{X}_I\vec{\beta}_I)'\mathbf{X}_{II}\vec{\beta}_{II} - \underbrace{(\mathbf{X}_I\vec{\beta}_I)'\mathbf{H}_1\mathbf{X}_{II}\vec{\beta}_{II}}_{\text{see below}} \\ &= (\mathbf{X}_I\vec{\beta}_I)'\mathbf{X}_{II}\vec{\beta}_{II} - (\mathbf{X}_I\vec{\beta}_I)'\mathbf{X}_{II}\vec{\beta}_{II} \\ &= \mathbf{O} \end{aligned}$$

since  $\{[(\mathbf{X}_I\vec{\beta}_I)'\mathbf{H}_1]\}' = \{\mathbf{H}_1\mathbf{X}_I\vec{\beta}_I\}' = (\mathbf{X}_I\vec{\beta}_I)'$ .

(3) A2Q5.

## ANOVA for Model Selection

$H_0: \vec{\beta}_{II} = \vec{0}$  versus  $H_A: \vec{\beta}_{II} \neq \vec{0}$ .

| Source                                       | df            | SS                  | MS                                 | Statistics |
|--|---------------|---------------------|------------------------------------|------------|
| Due to $\vec{\beta}$                         | $k$           | SSR(full)           | SSR(full)/ $k$                     |            |
| Due to $\vec{\beta}_I$                       | $\ell$        | SSR(reduced)        | SSR(reduced)/ $\ell$               | $F$        |
| Due to $\vec{\beta}_{II} \mid \vec{\beta}_I$ | $k - \ell$    | SSR(full   reduced) | SSR(full   reduced)/( $k - \ell$ ) |            |
| Error  | $n - (k + 1)$ | SSE                 | MSE                                |            |
| Total  | $n - 1$       | SST                 |                                    |            |

where

$$F = \frac{\text{SSR}(\text{full} \mid \text{reduced}) / (k - \ell)}{\text{SSE} / (n - k - 1)}$$

If we reject  $H_0$ , then the full model is better than the reduced model. Reject  $H_0$  when  $F > F_\alpha(k - \ell, n - k - 1)$ .



## Regression for Models without Full Rank

### DEFINITION 10.1: Generalized Inverse

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$ . The **generalized inverse** ( $g$ -inverse) of  $\mathbf{A}$  is any  $\mathbf{G} \in \mathbb{R}^{k \times n}$  satisfying

$$\mathbf{AGA} = \mathbf{A}.$$

We say  $\mathbf{G}$  is a  $g$ -inverse of  $\mathbf{A}$ .

### EXAMPLE 10.1

Let  $\mathbf{A} = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}$ . Find a  $g$ -inverse of  $\mathbf{A}$ .

**Solution:**

$$\begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} (a \quad b \quad c \quad d) \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}.$$

If we pick  $a = 1, b = c = d = 0$ , then  $\mathbf{G} = (1 \quad 0 \quad 0 \quad 0)$  is a  $g$ -inverse of  $\mathbf{A}$ . Also,  $\tilde{\mathbf{G}} = (1/2 \quad 1/4 \quad 0 \quad 0)$  is another  $g$ -inverse of  $\mathbf{A}$ . Hence, we can see that  $g$ -inverses are not unique.

### REMARK — Basic Facts

(1) If  $\mathbf{A}$  is invertible, then the  $g$ -inverse of  $\mathbf{A}$  is unique and given by  $\mathbf{A}^{-1}$ .

- **Proof:** Let  $\mathbf{G}$  be any  $g$ -inverse of  $\mathbf{A}$ , then  $\mathbf{AGA} = \mathbf{A}$ .

$$\mathbf{A}^{-1} \mathbf{A} \mathbf{G} \mathbf{A} \mathbf{A}^{-1} = \mathbf{A}^{-1} \mathbf{A} \mathbf{A}^{-1} \implies \mathbf{G} = \mathbf{A}^{-1}.$$

Clearly,  $\mathbf{A} \mathbf{A}^{-1} \mathbf{A} = \mathbf{A}$ .

(2) If  $\mathbf{G}$  is a  $g$ -inverse of  $\mathbf{A}$ , then for any  $\mathbf{C} \in \mathbb{R}^{k \times n}$ ,  $\mathbf{G}_1 = \mathbf{G} + \mathbf{C} - \mathbf{G} \mathbf{A} \mathbf{C} \mathbf{A} \mathbf{G}$  is also a  $g$ -inverse of  $\mathbf{A}$ .

- **Proof:** Note that

$$\begin{aligned} \mathbf{A} \mathbf{G}_1 \mathbf{A} &= \mathbf{A} \mathbf{G} \mathbf{A} + \mathbf{A} \mathbf{C} \mathbf{A} - \mathbf{A} \mathbf{G} \mathbf{A} \mathbf{C} \mathbf{A} \mathbf{G} \mathbf{A} \\ &= \mathbf{A} \mathbf{G} \mathbf{A} + \mathbf{A} \mathbf{C} \mathbf{A} - \mathbf{A} \mathbf{C} \mathbf{A} \\ &= \mathbf{A} \mathbf{G} \mathbf{A} \\ &= \mathbf{A}. \end{aligned}$$

### LEMMA 10.1

Every matrix  $\mathbf{A}$  has at least one  $g$ -inverse.

**Proof:** Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$  with  $\text{rank}(\mathbf{A}) = r < \min\{n, k\}$ . Then,

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix},$$

where  $\mathbf{A}_{11} \in \mathbb{R}^{r \times r}$  with  $\text{rank}(\mathbf{A}_{11}) = r$ .

Claim:

$$\mathbf{G} = \begin{pmatrix} \mathbf{A}_{11}^{-1} & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix}_{k \times n}$$

is a  $g$ -inverse of  $\mathbf{A}$ .

$$\begin{aligned} \mathbf{AGA} &= \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix} \begin{pmatrix} \mathbf{A}_{11}^{-1} & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{I}_r & \mathbf{O} \\ \mathbf{A}_{21}\mathbf{A}_{11}^{-1} & \mathbf{O} \end{pmatrix} \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12} \end{pmatrix}. \end{aligned}$$

Since  $\text{rank}(\mathbf{A}) = r = \text{rank}(\mathbf{A}_{11})$ , it follows that  $\mathbf{A}_{21}$  and  $\mathbf{A}_{22}$  are linear combinations of  $\mathbf{A}_{11}$  and  $\mathbf{A}_{12}$ . Thus, one can find a  $\mathbf{B} \in \mathbb{R}^{(n-r) \times r}$  such that

$$\begin{pmatrix} \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix} = \mathbf{B} \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \end{pmatrix}.$$

Hence,

$$\mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12} = \mathbf{B}\mathbf{A}_{11}\mathbf{A}_{11}^{-1}\mathbf{A}_{12} = \mathbf{B}\mathbf{A}_{12} = \mathbf{A}_{22}.$$

Therefore,  $\mathbf{AGA} = \mathbf{A}$ .

LECTURE 11  
13th February

## 11 Lecture 11: $g$ -inverse

### Algorithm for Finding a $g$ -inverse

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$  with  $\text{rank}(\mathbf{A}) = r < \min\{n, k\}$ .

- Step 1: Find an invertible sub-matrix  $\mathbf{M} \in \mathbb{R}^{r \times r}$ .
- Step 2: Compute  $(\mathbf{M}^{-1})'$ .
- Step 3: Replace  $\mathbf{M}$  with  $(\mathbf{M}^{-1})'$  in  $\mathbf{A}$ .
- Step 4: Set all other elements in  $\mathbf{A}$  to be 0.
- Step 5: Transpose the resulting matrix to  $\mathbf{G} \in \mathbb{R}^{k \times n}$ .

#### EXAMPLE 11.1

Compute a  $g$ -inverse of  $\mathbf{A} = \begin{pmatrix} 4 & 1 & 2 & 0 \\ 1 & 1 & 5 & 15 \\ 3 & 1 & 3 & 5 \end{pmatrix}$ .

**Solution:** Note that  $n = 3$  and  $k = 4$ . Let

$$\begin{aligned} \vec{v}_1 &= (4 \ 1 \ 2 \ 0), \\ \vec{v}_2 &= (1 \ 1 \ 5 \ 15), \\ \vec{v}_3 &= (3 \ 1 \ 3 \ 5). \end{aligned}$$

$\vec{v}_1$  and  $\vec{v}_2$  are linearly independent since

$$a\vec{v}_1 + b\vec{v}_2 = \vec{0} \implies 15b = 0 \text{ and } 4a = 0 \implies a = b = 0.$$

Also,  $3\vec{v}_3 = 2\vec{v}_1 + \vec{v}_2$ . Therefore,  $\text{rank}(\mathbf{A}) = 2$ . Now,

- Step 1:

$$\mathbf{M} = \begin{pmatrix} 4 & 0 \\ 3 & 5 \end{pmatrix}.$$

- Step 2:

$$(\mathbf{M}^{-1})' = \frac{1}{20} \begin{pmatrix} 5 & -3 \\ 0 & 4 \end{pmatrix} = \begin{pmatrix} 5/20 & -3/20 \\ 0 & 4/20 \end{pmatrix}.$$

- Step 3:

$$\begin{pmatrix} 5/20 & 1 & 2 & -3/20 \\ 1 & 1 & 5 & 15 \\ 0 & 1 & 3 & 4/20 \end{pmatrix}.$$

- Step 4:

$$\begin{pmatrix} 5/20 & 0 & 0 & -3/20 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 4/20 \end{pmatrix}.$$

- Step 5:

$$\begin{pmatrix} 5/20 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ -3/20 & 0 & 4/20 \end{pmatrix}.$$

Verify that  $\mathbf{AGA} = \mathbf{A}$ .

### THEOREM 11.1

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$  with  $\text{rank}(\mathbf{A}) = r < \min\{n, k\}$  and  $\mathbf{G}$  be a  $g$ -inverse of  $\mathbf{A}$ . Let  $\mathbf{F}$  be a  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ . Then,

- (1)  $\mathbf{G}'$  is a  $g$ -inverse of  $\mathbf{A}'$ .
- (2)  $\text{rank}(\mathbf{GA}) = \text{rank}(\mathbf{AG}) = \text{rank}(\mathbf{A}) = r$ .
- (3)  $\mathbf{A} = \mathbf{AFA}'\mathbf{A}$  and  $\mathbf{A}' = \mathbf{A}'\mathbf{AFA}'$ . This means that  $\mathbf{FA}'$  is a  $g$ -inverse of  $\mathbf{A}$ .

**Proof:**

- (1) Since  $\mathbf{AGA} = \mathbf{A}$ , we have that  $(\mathbf{AGA})' = \mathbf{A}'\mathbf{G}'\mathbf{A}' = \mathbf{A}'$ . Therefore,  $\mathbf{G}'$  is a  $g$ -inverse of  $\mathbf{A}'$ .
- (2) Since  $\mathbf{AGA} = \mathbf{A}$ ,  $\text{rank}(\mathbf{A}) \leq \text{rank}(\mathbf{GA}) \leq \text{rank}(\mathbf{A})$ . Similarly,  $\text{rank}(\mathbf{A}) \leq \text{rank}(\mathbf{AG}) \leq \text{rank}(\mathbf{A})$ . Therefore,  

$$\text{rank}(\mathbf{AG}) = \text{rank}(\mathbf{GA}) = \text{rank}(\mathbf{A}).$$
- (3) Since  $\mathbf{F}$  is a  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ , we have  $\mathbf{A}'\mathbf{AFA}'\mathbf{A} = \mathbf{A}'\mathbf{A}$ . Rearranging,

$$\begin{aligned} \mathbf{A}'\mathbf{AFA}'\mathbf{A} - \mathbf{A}'\mathbf{A} &= \mathbf{O} \\ (\mathbf{A}'\mathbf{AFA}' - \mathbf{A}')\mathbf{A} &= \mathbf{O} \\ \mathbf{A}'(\mathbf{AFA}'\mathbf{A} - \mathbf{A}) &= \mathbf{O}. \end{aligned}$$

Note that  $(\mathbf{AFA}'\mathbf{A})' = \mathbf{A}'\mathbf{AF}'\mathbf{A}$  and

$$\begin{aligned}\mathbf{A}'\mathbf{AF}'\mathbf{A}'(\mathbf{AFA}'\mathbf{A} - \mathbf{A}) &= \mathbf{A}'\mathbf{AF}'\underbrace{\mathbf{A}'\mathbf{AFA}'\mathbf{A}}_{\mathbf{A}'\mathbf{A}} - \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A} \\ &= \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A} - \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A} \\ &= \mathbf{O}.\end{aligned}$$

Therefore,

$$(\mathbf{AFA}'\mathbf{A} - \mathbf{A})'(\mathbf{AFA}'\mathbf{A} - \mathbf{A}) = \mathbf{O}.$$

Hence,

$$\mathbf{AFA}'\mathbf{A} - \mathbf{A} = \mathbf{O} \implies \mathbf{AFA}'\mathbf{A} = \mathbf{A}.$$

Similarly,  $\mathbf{A}'\mathbf{AFA}'\mathbf{A} = \mathbf{A}'\mathbf{A}$ , which implies

$$(\mathbf{A}'\mathbf{AFA}' - \mathbf{A}')\mathbf{A} = \mathbf{O}.$$

By direct calculation,

$$(\mathbf{A}'\mathbf{AFA}' - \mathbf{A}')\mathbf{AF}'\mathbf{A}'\mathbf{A} = \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A} - \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A} = \mathbf{O}.$$

Therefore,

$$(\mathbf{A}'\mathbf{AFA}' - \mathbf{A}')(\mathbf{A}'\mathbf{AFA}' - \mathbf{A}')' = \mathbf{O}.$$

Hence,

$$\mathbf{A}'\mathbf{AFA}' = \mathbf{A}'.$$

$\mathbf{AF}$  is a  $g$ -inverse of  $\mathbf{A}'$  and  $\mathbf{FA}'$  is a  $g$ -inverse of  $\mathbf{A}$ .

### THEOREM 11.2

Let  $\mathbf{F}$  be a  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ .

- (1)  $\mathbf{F}'$  is a  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ .
- (2)  $\text{rank}(\mathbf{AFA}') = \text{rank}(\mathbf{A})$ .
- (3) Let  $\tilde{\mathbf{F}}$  be any  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ , then  $\mathbf{A}'\mathbf{FA} = \mathbf{A}'\tilde{\mathbf{F}}\mathbf{A}$ .
- (4)  $\mathbf{AFA}'$  is symmetric.

**Proof:**

- (1) Using Theorem 11.1,  $\mathbf{A}'\mathbf{AFA}'\mathbf{A} = \mathbf{A}'\mathbf{A}$ , so

$$\mathbf{A}'\mathbf{A} = (\mathbf{A}'\mathbf{A})' = \mathbf{A}'\mathbf{AF}'\mathbf{A}'\mathbf{A}.$$

- (2) By Theorem 11.1, we have  $\mathbf{A} = \mathbf{AFA}'\mathbf{A}$ . It follows that

$$\text{rank}(\mathbf{A}) \leq \text{rank}(\mathbf{AFA}') \leq \text{rank}(\mathbf{A}).$$

Therefore,  $\text{rank}(\mathbf{AFA}') = \text{rank}(\mathbf{A})$ .

- (3) Let  $\tilde{\mathbf{F}}$  be any  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ . Then,  $\mathbf{A} = \mathbf{AFA}'\mathbf{A} = \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}'\mathbf{A}$  by Theorem 11.1, so

$$(\mathbf{AFA}' - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}')\mathbf{A} = \mathbf{O}.$$

Therefore,

$$\begin{aligned}
(\mathbf{AFA}' - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}')(\mathbf{AFA}' - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}')' &= (\mathbf{AFA}' - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}')(\mathbf{AF}'\mathbf{A}' - \mathbf{A}\tilde{\mathbf{F}}'\mathbf{A}')' \\
&= (\mathbf{AFA}' - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}')\mathbf{A}(\mathbf{F}'\mathbf{A}' - \tilde{\mathbf{F}}'\mathbf{A}') \\
&= \underbrace{(\mathbf{AFA}'\mathbf{A} - \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}'\mathbf{A})}_{\mathbf{A}}(\mathbf{F}'\mathbf{A}' - \tilde{\mathbf{F}}'\mathbf{A}') \\
&= \mathbf{O}.
\end{aligned}$$

Hence,  $\mathbf{AFA}' = \mathbf{A}\tilde{\mathbf{F}}\mathbf{A}'$ .

(4) By (1),  $\mathbf{F}'$  is a  $g$ -inverse of  $\mathbf{A}'\mathbf{A}$ . Hence,  $\mathbf{AFA}' = \mathbf{AF}'\mathbf{A}' = (\mathbf{AFA}')'$ . Therefore,  $\mathbf{AFA}'$  is symmetric.

### THEOREM 11.3

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$ . Consider the system of equations

$$\mathbf{A}\vec{x} = \vec{y}.$$

(1) If  $\vec{x}_0$  is a solution of the system of equations, then  $\mathbf{GA}\vec{x}_0$  is also a solution of the system of equations for any  $g$ -inverse  $\mathbf{G}$  of  $\mathbf{A}$ .

(2) Let  $\mathbf{G}$  be a  $g$ -inverse of  $\mathbf{A}$ , then for any  $\vec{z} \in \mathbb{R}^k$ ,

$$\mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}$$

is a solution of the system of equations.

(3) Every solution can be written in the form of (2).

**Proof:**

(1)  $\vec{x}_0$  is a solution implies that  $\mathbf{A}\vec{x}_0 = \vec{y}$ . However,

$$\mathbf{A}(\mathbf{GA}\vec{x}_0) = (\mathbf{AGA})\vec{x}_0 = \mathbf{A}\vec{x}_0 = \vec{y}.$$

(2) Note that

$$\{\mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z} \mid \vec{z} \in \mathbb{R}^k\}.$$

So,

$$\begin{aligned}
\mathbf{A}(\mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}) &= \mathbf{AG}\vec{y} + (\mathbf{AGA}\vec{z} - \mathbf{A}\vec{z}) \\
&= \mathbf{AG}\vec{y} \\
&= \mathbf{AG}(\mathbf{A}\vec{x}) \\
&= \mathbf{AGA}\vec{x} \\
&= \mathbf{A}\vec{x} \\
&= \vec{y}.
\end{aligned}$$

(3) Let  $\vec{x}_0$  be any solution, so  $\mathbf{A}\vec{x}_0 = \vec{y}$ . Choose  $\vec{z} = (\mathbf{GA} - \mathbf{I})\vec{x}_0$ .

$$\begin{aligned}
\mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z} &= \mathbf{G}\vec{y} + (\mathbf{GAGA} - 2\mathbf{GA} + \mathbf{I})\vec{x}_0 \\
&= \mathbf{G}\vec{y} + (-\mathbf{GA} + \mathbf{I})\vec{x}_0 \\
&= \mathbf{G}\vec{y} - \mathbf{GA}\vec{x}_0 + \vec{x}_0 \\
&= \mathbf{GA}\vec{x}_0 - \mathbf{GA}\vec{x}_0 + \vec{x}_0 \\
&= \vec{x}_0.
\end{aligned}$$

## 12 Lecture 12: Regression Without Full Rank

### DEFINITION 12.1: Estimable

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$  with  $\text{rank}(\mathbf{A}) = r < \min\{n, k\}$ ,  $\vec{x} \in \mathbb{R}^k$ . Given a  $\vec{b} \in \mathbb{R}^{k \times 1}$ , the quantity  $\vec{b}'\vec{x}$  is called **estimable** if its value is the same for every solution of  $\mathbf{A}\vec{x} = \vec{y}$ .

### THEOREM 12.1

Let  $\mathbf{A} \in \mathbb{R}^{n \times k}$ ,  $\vec{b} \in \mathbb{R}^k$ , and  $\mathbf{G}$  be a  $g$ -inverse of  $\mathbf{A}$ .

$$\vec{b}'\vec{x} \text{ is estimable} \iff \vec{b}'\mathbf{GA} = \vec{b}'.$$

**Proof:** Let  $\vec{x}_1$  and  $\vec{x}_2$  be any two solutions of  $\mathbf{A}\vec{x} = \vec{y}$ . By Theorem 11.3, there exists  $\vec{z}_1, \vec{z}_2 \in \mathbb{R}^k$  such that

$$\begin{aligned}\vec{x}_1 &= \mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}_1 \\ \vec{x}_2 &= \mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}_2.\end{aligned}$$

( $\Leftarrow$ ) *Sufficiency:* Assume that  $\vec{b}'\mathbf{GA} = \vec{b}'$ .

$$\begin{aligned}\implies \vec{b}'\mathbf{GA} - \vec{b}' &= \vec{0}' \\ \implies \vec{b}'(\mathbf{GA} - \mathbf{I}) &= \vec{0}' \\ \implies \vec{b}'(\mathbf{GA} - \mathbf{I})\vec{z} &= 0\end{aligned}$$

Now,

$$\begin{aligned}\vec{b}'\vec{x}_1 - \vec{b}'\vec{x}_2 &= \vec{b}'(\vec{x}_1 - \vec{x}_2) \\ &= \vec{b}'[(\mathbf{GA} - \mathbf{I})(\vec{z}_1 - \vec{z}_2)] \\ &= \vec{b}'(\mathbf{GA} - \mathbf{I})(\vec{z}_1 - \vec{z}_2) \\ &= 0.\end{aligned}$$

Hence,  $\vec{b}'\vec{x}_1 = \vec{b}'\vec{x}_2 \implies \vec{b}'\vec{x}$  is estimable.

( $\implies$ ) *Necessity:* Assume that  $\vec{b}'\vec{x}$  is estimable. Let  $\vec{x}_0$  be a solution of  $\mathbf{A}\vec{x} = \vec{y}$ . Choose  $\vec{z}_0$  such that

$$\vec{x}_0 = \mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}_0.$$

Let  $z \in \mathbb{R}^k$ . By definition,

$$\vec{x} = \mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})(\vec{z}_0 - \vec{z}).$$

Hence,  $\vec{x}$  is a solution of  $\mathbf{A}\vec{x} = \vec{y}$ , and so

$$\begin{aligned}\vec{b}'\vec{x} &= \vec{b}'[\mathbf{G}\vec{y} + (\mathbf{GA} - \mathbf{I})\vec{z}_0 + (\mathbf{GA} - \mathbf{I})\vec{z}] \\ &= \vec{b}'\vec{x}_0 + \vec{b}'(\mathbf{GA} - \mathbf{I})\vec{z}.\end{aligned}$$

So,  $\vec{b}'(\mathbf{GA} - \mathbf{I})\vec{z} = 0$  for all  $\vec{z}$ , therefore  $\vec{b}'(\mathbf{GA} - \mathbf{I}) = \vec{0}' \implies \vec{b}'\mathbf{GA} = \vec{b}'$ .

## Non-Full Rank Regression

In regression,  $\vec{Y} = \mathbf{X}\vec{\beta} + \vec{\varepsilon}$  and our normal equation is  $\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}$ . We consider the case where  $\mathbf{X}'\mathbf{X}$  is not invertible. Let  $\mathbf{F}$  be a  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$ , so

$$\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y}.$$

Claim:  $\vec{\beta}_0$  is a solution of

$$\mathbf{X}'\mathbf{X}\vec{\beta}_0 = \mathbf{X}'\vec{Y}.$$

By Theorem 11.1,

$$\mathbf{X}'\mathbf{X}\vec{\beta}_0 = \mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\vec{Y} = \mathbf{X}'\vec{Y}.$$

However,  $\vec{\beta}_0$  may not be a good estimator.

### REMARK — Properties of $\vec{\beta}_0$

(1) Expectation:  $\mathbb{E}[\vec{\beta}_0] = \mathbb{E}[\mathbf{F}\mathbf{X}'\vec{Y}] = \mathbf{F}\mathbf{X}'\mathbb{E}[\vec{Y}] = \mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta} \neq \vec{\beta}$  in general.

(2) Variance:

$$\begin{aligned} \text{Var}(\vec{\beta}_0) &= \text{Cov}(\mathbf{F}\mathbf{X}'\vec{Y}, \mathbf{F}\mathbf{X}'\vec{Y}) \\ &= \mathbf{F}\mathbf{X}'\text{Cov}(\vec{Y}, \vec{Y})\mathbf{X}\mathbf{F}' \\ &= \sigma^2\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}'. \end{aligned}$$

(3) Fitted values:

$$\begin{aligned} \hat{\vec{Y}} &= \mathbf{X}\vec{\beta}_0 \\ &= \mathbf{X}\mathbf{F}\mathbf{X}'\vec{Y} \\ &= (\mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}. \end{aligned}$$

Note that by Theorem 11.2 (3),  $\hat{\vec{Y}}$  does not depend on which  $g$ -inverse ( $\mathbf{F}$ ) we use.

(4) SSE:

$$\begin{aligned} \text{SSE} &= (\vec{Y} - \hat{\vec{Y}})'(\vec{Y} - \hat{\vec{Y}}) \\ &= (\vec{Y} - \mathbf{X}\mathbf{F}\mathbf{X}'\vec{Y})'(\vec{Y} - \mathbf{X}\mathbf{F}\mathbf{X}'\vec{Y}) \\ &= \vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y} \\ &= \vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y} && \text{Theorem 11.2 (4)} \\ &= \vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}, \end{aligned}$$

where

$$\begin{aligned} (\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}') &= \mathbf{I} - 2\mathbf{X}\mathbf{F}\mathbf{X}' + \underbrace{\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'}_{\mathbf{X}} \\ &= \mathbf{I} - 2\mathbf{X}\mathbf{F}\mathbf{X}' + \mathbf{X}\mathbf{F}\mathbf{X}' && \text{Theorem 11.1 (3)} \\ &= \mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}'. \end{aligned}$$

**EXAMPLE 12.1**

| Weights of Six Plants |          |          |
|-----------------------|----------|----------|
| Types of Plants       |          |          |
| Normal                | Off-type | Aberrant |
| 101                   | 84       | 32       |
| 105                   | 88       |          |
| 94                    |          |          |
| 300                   | 172      | 32       |

Relation between weight and types. Let  $Y = \text{weight}$ ,  $x_1 = \text{normal}$ ,  $x_2 = \text{off-type}$ ,  $x_3 = \text{aberrant}$ , where all the covariates are binary.  $Y_{ij}$  = observation of  $j^{\text{th}}$  plant of type  $i$  for  $i = 1, 2, 3$ ;  $n_1 = 3$ ,  $n_2 = 2$ ,  $n_3 = 1$ , and  $n = n_1 + n_2 + n_3 = 6$ .

$$\vec{Y}' = (Y_{11} \ Y_{12} \ Y_{13} \ Y_{21} \ Y_{22} \ Y_{33}) = (101 \ 105 \ 94 \ 84 \ 88 \ 32)'.$$

Regression model:

$$Y_{ij} = \beta_0 + \beta_i + \varepsilon_{ij}.$$

- $\beta_0$  = population mean;
- $\beta_i$  = effect of type  $i$  on the weight;
- $\varepsilon_{ij}$  = random error of observation  $Y_{ij}$ .

Explicitly, we have

$$\begin{aligned} Y_{11} &= \beta_0 + \beta_1 + 0\beta_2 + 0\beta_3 + \varepsilon_{11} \\ Y_{12} &= \beta_0 + \beta_1 + 0\beta_2 + 0\beta_3 + \varepsilon_{12} \\ Y_{13} &= \beta_0 + \beta_1 + 0\beta_2 + 0\beta_3 + \varepsilon_{13} \\ Y_{21} &= \beta_0 + 0\beta_1 + \beta_2 + 0\beta_3 + \varepsilon_{21} \\ Y_{22} &= \beta_0 + 0\beta_1 + \beta_2 + 0\beta_3 + \varepsilon_{22} \\ Y_{33} &= \beta_0 + 0\beta_1 + 0\beta_2 + \beta_3 + \varepsilon_{33}. \end{aligned}$$

Therefore,  $\vec{Y} = \mathbf{X}\vec{\beta}$  with

$$\mathbf{X} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

Note that  $\text{rank}(\mathbf{X}) = 3 < \min\{6, 4\}$  so  $\mathbf{X}$  is not full rank.

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} 6 & 3 & 2 & 1 \\ 3 & 3 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

$$\mathbf{X}'\vec{Y} = \begin{pmatrix} Y_{..} \\ Y_{1.} \\ Y_{2.} \\ Y_{3.} \end{pmatrix} = \begin{pmatrix} 504 \\ 300 \\ 172 \\ 32 \end{pmatrix}.$$



Normal equation  $\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}$ .  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$ :

$$\mathbf{M} = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \implies \mathbf{F} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1/3 & 0 & 0 \\ 0 & 0 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Note that

$$\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y} = \begin{pmatrix} 0 \\ 100 \\ 86 \\ 32 \end{pmatrix}$$

is one solution. However, we cannot claim that  $\vec{\beta}_0$  is an estimate of  $\vec{\beta}$ . By direct calculation,

$$\mathbf{F}\mathbf{X}'\mathbf{X} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1/3 & 0 & 0 \\ 0 & 0 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 6 & 3 & 2 & 1 \\ 3 & 3 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

Choose  $\vec{b}' = (1 \ 1 \ 0 \ 0)$  and compute

$$\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = (1 \ 1 \ 0 \ 0) = \vec{b}',$$

so  $\vec{b}'\vec{\beta} = \beta_0 + \beta_1$  is estimable. Therefore,  $\hat{\beta}_0 + \hat{\beta}_1$  is an estimator of  $\beta_0 + \beta_1$ .

LECTURE 13  
27th February

## 13 Lecture 13: Regression Without Full Rank (Continued)

### THEOREM 13.1

If  $\text{rank}(\mathbf{X}) = r$ , then

$$S^2 = \hat{\sigma}^2 = \frac{\text{SSE}}{n - r}$$

is an unbiased estimator of  $\sigma^2$ .

**Proof:**

$$\begin{aligned} \mathbb{E}[\text{SSE}] &= \mathbb{E}[\vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}] \\ &= \text{tr}((\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\sigma^2\mathbf{I}) + (\mathbf{X}\vec{\beta})'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\mathbf{X}\vec{\beta} \\ &= \sigma^2(\text{tr}(\mathbf{I}) - \text{tr}(\mathbf{X}\mathbf{F}\mathbf{X}')) + (\mathbf{X}\vec{\beta})'\mathbf{X}\vec{\beta} - \vec{\beta}'\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta} \\ &= \sigma^2(n - r) + (\mathbf{X}\vec{\beta})'\mathbf{X}\vec{\beta} - (\mathbf{X}\vec{\beta})'(\mathbf{X}\vec{\beta}) \\ &= \sigma^2(n - r), \end{aligned}$$

where we used the fact that  $\text{rank}(\mathbf{X}\mathbf{F}\mathbf{X}') = \text{rank}(\mathbf{X}) = r$  by Theorem 11.2 (2). Therefore,

$$\mathbb{E}[\text{SSE}] = \sigma^2(n - r) \implies \mathbb{E}[\hat{\sigma}^2] = \sigma^2.$$

**THEOREM 13.2**

If  $\text{rank}(\mathbf{X}) = r$  and  $\vec{Y} \sim MN(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I})$ , then

- (1)  $\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y} \sim MN(\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}, \sigma^2\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}')$ .
- (2)  $\vec{\beta}_0$  and  $\hat{\sigma}^2$  are independent.
- (3)  $SSE/\sigma^2 \sim \chi^2(n - r)$ .
- (4)  $SSR/\sigma^2 \sim \chi^2(r - 1, \lambda)$ , where

$$\lambda = \frac{1}{2\sigma^2}(\mathbf{X}\vec{\beta})'(\mathbf{X}\mathbf{F}\mathbf{X}' - \frac{1}{n}\mathbf{J})\mathbf{X}\vec{\beta}.$$

- (5)  $SSE$  and  $SSR$  are independent.

**Proof:**

- (1) Trivial.

- (2)  $\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y}$  and

$$\hat{\sigma}^2 = \frac{SSE}{n - r} = \frac{1}{n - r}\vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}.$$

By direct calculation,

$$\begin{aligned} & \mathbf{F}\mathbf{X}'\sigma^2\mathbf{I}(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}') \\ &= \frac{\sigma^2}{n - r}\mathbf{F}\mathbf{X}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}') \\ &= \frac{\sigma^2}{n - r}[\mathbf{F}\mathbf{X}' - \mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'] \\ &= \frac{\sigma^2}{n - r}[\mathbf{F}\mathbf{X}' - \mathbf{F}\mathbf{X}'] \quad \text{Theorem 11.1} \\ &= \mathbf{O}. \end{aligned}$$

Therefore,  $\vec{\beta}_0$  and  $\hat{\sigma}^2$  are independent by Theorem 5.1.

- (3) Note that

$$\frac{SSE}{\sigma^2} = \frac{\vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}}{\sigma}$$

and

$$\frac{\vec{Y}}{\sigma} \sim MN\left(\frac{\mathbf{X}\vec{\beta}}{\sigma}, \mathbf{I}\right).$$

Note that  $\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}'$  is idempotent (see properties of  $\vec{\beta}_0$ ). Now,

$$\begin{aligned} \lambda &= \frac{1}{2}\left(\frac{\mathbf{X}\vec{\beta}}{\sigma}\right)'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\frac{\mathbf{X}\vec{\beta}}{\sigma} \\ &= \frac{1}{2\sigma^2}\vec{\beta}'[\mathbf{X}'\mathbf{X} - \mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X}]\vec{\beta} \\ &= \frac{1}{2\sigma^2}\vec{\beta}'[\mathbf{X}'\mathbf{X} - \mathbf{X}'\mathbf{X}]\vec{\beta} \\ &= 0. \end{aligned}$$

The result follows from Theorem 4.1.

(4) Note that

$$\frac{SSR}{\sigma^2} = \frac{\vec{Y}'}{\sigma} (\mathbf{XFX}' - \frac{1}{n}\mathbf{J}) \frac{\vec{Y}}{\sigma}.$$

We need to show that  $\mathbf{XFX}' - \frac{1}{n}\mathbf{J}$  is idempotent.

$$\begin{aligned} (\mathbf{XFX}' - \frac{1}{n}\mathbf{J})^2 &= (\mathbf{XFX}' - \frac{1}{n}\mathbf{J})(\mathbf{XFX}' - \frac{1}{n}\mathbf{J}) \\ &= \mathbf{XFX}'\mathbf{XFX}' - \frac{1}{n}\mathbf{JXFX}' - \frac{1}{n}\mathbf{XFX}'\mathbf{J} + \frac{1}{n^2}\mathbf{J}\mathbf{J} \\ &= \mathbf{XFX}' - \frac{1}{n}\mathbf{XFX}'\mathbf{J} - \frac{1}{n}\mathbf{JXFX}' + \frac{1}{n}\mathbf{J}. \end{aligned}$$

We know that  $\mathbf{XFX}'\mathbf{X} = \mathbf{X}$ , so partitioning we see

$$\mathbf{XFX}' \begin{pmatrix} \vec{j} \\ \mathbf{X}_1 \end{pmatrix} = \mathbf{X},$$

which implies that  $\mathbf{XFX}'\vec{j} = \vec{j}$ . Therefore,  $\mathbf{XFX}'\mathbf{J} = \mathbf{J}$ . Continuing,

$$\begin{aligned} \mathbf{XFX}' - \frac{1}{n}\mathbf{XFX}'\mathbf{J} - \frac{1}{n}\mathbf{JXFX}' + \frac{1}{n}\mathbf{J} &= \mathbf{XFX}' - \frac{1}{n}\mathbf{J} - \frac{1}{n}\mathbf{J} + \frac{1}{n}\mathbf{J} \\ &= \mathbf{XFX}' - \frac{1}{n}\mathbf{J}, \end{aligned}$$

so  $\mathbf{XFX}'$  is idempotent. The result follows from Theorem 4.1.

(5)  $SSE = \vec{Y}'(\mathbf{I} - \mathbf{XFX}')\vec{Y}$  and  $SSR = \vec{Y}'(\mathbf{XFX}' - \frac{1}{n}\mathbf{J})\vec{Y}$ .

$$\begin{aligned} (\mathbf{I} - \mathbf{XFX}')(\mathbf{XFX}' - \frac{1}{n}\mathbf{J}) &= \mathbf{XFX}' - \frac{1}{n}\mathbf{J} - \mathbf{XFX}'\mathbf{XFX}' + \frac{1}{n}\mathbf{XFX}'\mathbf{J} \\ &= \mathbf{XFX}' - \frac{1}{n}\mathbf{J} - \mathbf{XFX}' + \frac{1}{n}\mathbf{J} \\ &= \mathbf{O}. \end{aligned}$$

The result follows from Theorem 5.2.

## ANOVA Table

| Source of Variation  | Degrees of Freedom | Sum of Squares | Mean Square         | $F$       |
|----------------------|--------------------|----------------|---------------------|-----------|
| Due to $\vec{\beta}$ | $r$                | SSR            | $MSR = SSR/(r - 1)$ | $MSR/MSE$ |
| Error                | $n - r$            | SSE            | $MSE = SSE/(n - r)$ |           |
| Total                | $n - 1$            | SST            |                     |           |

Rejection region:  $F > F_\alpha(r - 1, n - r)$ , and we are testing  $H_0: \mathbf{X}\vec{\beta} = \vec{0}$  versus  $H_A: \mathbf{X}\vec{\beta} \neq \vec{0}$ , which is not the same as the hypothesis test of  $H_0: \vec{\beta} = 0$  versus  $H_A: \vec{\beta} \neq 0$  as before.

### THEOREM 13.3

Let  $\vec{\beta} = \begin{pmatrix} \beta_0 \\ \vdots \\ \beta_k \end{pmatrix} \in \mathbb{R}^{k+1}$ ,  $\vec{b} \in \mathbb{R}^{k+1}$ , and  $\mathbf{F}$  be a  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$ .

$\vec{b}'\vec{\beta}$  is estimable if and only if one of the following hold:

- (1)  $\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}'$ .
- (2) There exists  $\vec{a} \in \mathbb{R}^n$  such that  $\vec{b}' = \vec{a}'\mathbf{X}$ .
- (3) There exists  $\vec{c} \in \mathbb{R}^{k+1}$  such that

$$\vec{b}' = \vec{c}'\mathbf{X}'\mathbf{X}.$$

**Proof:**

(1) Theorem 12.1.

(2) To show (1)  $\implies$  (2), choose  $\vec{a}' = \vec{b}'\mathbf{F}\mathbf{X}'$ , which implies that  $\vec{a}'\mathbf{X} = \vec{b}'$ . To show (2)  $\implies$  (1), note that  $\vec{b}' = \vec{a}'\mathbf{X}$ , and multiply to get  $\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{a}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{a}'\mathbf{X}$ .

(3) To show (1)  $\implies$  (3), choose  $\vec{c}' = \vec{b}'\mathbf{F}$ . To show (3)  $\implies$  (1),

$$\begin{aligned}\vec{b}' &= \vec{c}'\mathbf{X}'\mathbf{X} \\ \vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} &= \vec{c}'\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X} \\ &= \vec{c}'\mathbf{X}'\mathbf{X} \\ &= \vec{b}'.\end{aligned}$$

**REMARK**

Assume that  $\vec{b}'\vec{\beta}$  is estimable. Let  $\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y}$ .  $\vec{b}'\vec{\beta}_0$  is an estimator of  $\vec{b}'\vec{\beta}$ .

The expectation of  $\vec{b}'\vec{\beta}_0$  is

$$\begin{aligned}\mathbb{E}[\vec{b}'\vec{\beta}_0] &= \vec{b}'\mathbb{E}[\vec{\beta}_0] \\ &= \vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta} && \text{Theorem 13.2 (1)} \\ &= \vec{b}'\vec{\beta} && \text{Theorem 13.3 (1)}.\end{aligned}$$

Hence,  $\vec{b}'\vec{\beta}_0$  is an unbiased estimator of  $\vec{b}'\vec{\beta}$ .  
The variance of  $\vec{b}'\vec{\beta}_0$  is

$$\begin{aligned}\text{Var}(\vec{b}'\vec{\beta}_0) &= \vec{b}'\text{Var}(\vec{\beta}_0)\vec{b} \\ &= \vec{b}'\sigma^2\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}'\vec{b} && \text{Theorem 13.2 (1)} \\ &= \sigma^2\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X}'\mathbf{F}'\vec{b} \\ &= \sigma^2\vec{b}'\mathbf{F}'\vec{b} && \text{Theorem 13.3 (1)} \\ &= \sigma^2\vec{b}'\mathbf{F}\vec{b} && \text{Theorem 11.2 (3)}\end{aligned}$$

by Theorem 11.2 (3).

**THEOREM 13.4**

If  $\vec{Y} \sim MN(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I})$ , then

$$\vec{b}'\vec{\beta}_0 \sim MN(\vec{b}'\vec{\beta}, \sigma^2\vec{b}'\mathbf{F}\vec{b}).$$

Claim:

$$\frac{\vec{b}'\vec{\beta}_0 - \vec{b}'\vec{\beta}}{\sqrt{\sigma^2\vec{b}'\mathbf{F}\vec{b}}} \sim t_{n-r}.$$

$$\begin{aligned} \frac{\vec{b}'\vec{\beta}_0 - \vec{b}'\vec{\beta}}{\sqrt{\sigma^2 \vec{b}'\mathbf{F}\vec{b}}} &= \frac{\vec{b}'\vec{\beta}_0 - \vec{b}'\vec{\beta}}{\sqrt{\sigma^2 \vec{b}'\mathbf{F}\vec{b} \frac{\hat{\sigma}^2}{\sigma^2}}} \\ &= \frac{\vec{b}'\vec{\beta}_0 - \vec{b}'\vec{\beta}}{\sqrt{\sigma^2 \vec{b}'\mathbf{F}\vec{b}}} \bigg/ \sqrt{\frac{\hat{\sigma}^2}{\sigma^2}} \\ &= \frac{Z}{\sqrt{\frac{V}{n-r}}} \sim t(n-r), \end{aligned}$$

since

$$\frac{\hat{\sigma}^2}{\sigma^2} = \frac{\text{SSE}}{n-r},$$

$\text{SSE} \sim \chi^2(n-r)$ ,  $Z \sim \mathcal{N}(0,1)$ , and  $V \sim \chi^2(n-r)$ . By Theorem 13.2 (2)  $Z$  and  $V$  are independent.

A  $(1-\alpha)100\%$  confidence interval for  $\vec{b}'\vec{\beta}$  is given by

$$\vec{b}'\vec{\beta}_0 \pm t_{n-r;\alpha/2} \hat{\sigma} \sqrt{\vec{b}'\mathbf{F}\vec{b}}.$$

### EXAMPLE 13.1

Refer to Example 12.1. Note that

$$\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}' \implies b_2 + b_3 + b_4 = b_1, \quad b_2 = -b_3 = 1, \quad \vec{b}' = (0 \quad 1 \quad -1 \quad 0).$$

We can estimate

$$\vec{b}'\vec{\beta} = (0 \quad 1 \quad -1 \quad 0) \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \beta_1 - \beta_2.$$

$n = 6$ ,  $r = 3$ ,  $\alpha = 5\%$ ,  $t_{n-r;\alpha/2} = 3.182$ .

$$\begin{aligned} \hat{\sigma}^2 &= \frac{\text{SSE}}{n-r} = \frac{70}{3} = 23.33 \\ \frac{\vec{b}'\mathbf{F}\vec{b}}{\sigma^2} &= \frac{5}{6}. \end{aligned}$$

The confidence interval is

$$\vec{b}'\vec{\beta}_0 \pm 3.182 \sqrt{23.33} \sqrt{\frac{5}{6}} = 14 \pm 14.0303 = (-0.0303, 28.0303).$$

LECTURE 14  
2nd March

## 14 Lecture 14: General Linear Hypothesis Testing

Let  $\mathbf{B} \in \mathbb{R}^{(k+1) \times s}$  with  $\text{rank}(\mathbf{B}) = s$  and  $\vec{m} \in \mathbb{R}^s$ . We want to test  $H_0: \mathbf{B}'\vec{\beta} = \vec{m}$  versus  $H_A: \mathbf{B}'\vec{\beta} \neq \vec{m}$ . Write

$$\mathbf{B} = \begin{pmatrix} b_{11} & \cdots & b_{1s} \\ b_{21} & \cdots & b_{2s} \\ \vdots & \ddots & \vdots \\ b_{(k+1)1} & \cdots & b_{(k+1)s} \end{pmatrix} = (\vec{b}_1 \quad \vec{b}_2 \quad \cdots \quad \vec{b}_s).$$

Hence,

$$\mathbf{B}'\vec{\beta} = (\vec{b}_1'\vec{\beta} \quad \cdots \quad \vec{b}_s'\vec{\beta})'.$$

In order for  $\mathbf{B}'\vec{\beta}$  to be estimable, we need  $\vec{b}'_i\vec{\beta}$  to be estimable for  $i = 1, \dots, s$ . This requires

$$\vec{b}'_i\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}'_i \iff \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X} = \mathbf{B}'.$$

#### THEOREM 14.1

Let  $\vec{\beta}_0$  be a solution to the normal equation

$$\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}.$$

Since  $\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X} = \mathbf{X}$ , it follows that  $\mathbf{X}\vec{\beta}$  is estimable and  $\mathbf{X}\vec{\beta}_0$  is an estimator of  $\mathbf{X}\vec{\beta}$ . Assume that  $\mathbf{B}'\vec{\beta}$  is estimable.

(1)  $\mathbb{E}[\mathbf{B}'\vec{\beta}_0] = \mathbf{B}'\vec{\beta}.$

(2)  $\text{Var}(\mathbf{B}'\vec{\beta}_0) = \sigma^2\mathbf{B}'\mathbf{F}\mathbf{B}.$

**Proof:** Since  $\mathbf{B}'\vec{\beta}$  is estimable,  $\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X} = \mathbf{B}'$ . Furthermore,

$$\mathbf{B}'\vec{\beta}_0 = \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}_0.$$

(1) For the expectation,

$$\begin{aligned} \mathbb{E}[\mathbf{B}'\vec{\beta}_0] &= \mathbb{E}[\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}_0] \\ &= \mathbf{B}'\mathbf{F}\mathbb{E}[\mathbf{X}'\mathbf{X}\vec{\beta}_0] \\ &= \mathbf{B}'\mathbf{F}\mathbb{E}[\mathbf{X}'\vec{Y}] \\ &= \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbb{E}[\vec{Y}] \\ &= \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta} \\ &= \mathbf{B}'\vec{\beta}. \end{aligned}$$

(2) For the variance,

$$\begin{aligned} \text{Var}(\mathbf{B}'\vec{\beta}_0) &= \text{Var}(\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}_0) \\ &= \text{Var}(\mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y}) \\ &= (\mathbf{B}'\mathbf{F}\mathbf{X}')\sigma^2\mathbf{I}(\mathbf{B}'\mathbf{F}\mathbf{X}')' \\ &= \sigma^2\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}'\mathbf{B} \\ &= \sigma^2\mathbf{B}'\mathbf{F}\mathbf{B}. \end{aligned}$$

#### THEOREM 14.2

If  $\mathbf{B}'\vec{\beta}$  is estimable, then  $\text{rank}(\mathbf{B}'\mathbf{F}\mathbf{B}) = s$ .

**Proof:**  $\mathbf{B}'\vec{\beta}$  is estimable implies that there exists  $\mathbf{C} \in \mathbb{R}^{(k+1) \times s}$  matrix such that  $\mathbf{B}' = \mathbf{C}'\mathbf{X}'\mathbf{X}$  (extension of Theorem 13.3). We know that  $\text{rank}(\mathbf{B}') = s = \text{rank}(\mathbf{C}'\mathbf{X}'\mathbf{X}) \leq \text{rank}(\mathbf{C}') \leq s$ . Therefore,  $\text{rank}(\mathbf{C}) = \text{rank}(\mathbf{C}') = s$ . Also,  $\text{rank}(\mathbf{B}') \leq \text{rank}(\mathbf{C}'\mathbf{X}) \leq \text{rank}(\mathbf{C}') = s$ , which implies that  $\text{rank}(\mathbf{C}'\mathbf{X}') = \text{rank}(\mathbf{X}\mathbf{C}) = s$ . Now,

$$\begin{aligned} \mathbf{B}'\mathbf{F}\mathbf{B} &= \mathbf{C}'\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{C} \\ &= \mathbf{C}'\mathbf{X}'\mathbf{X}\mathbf{C} \\ &= (\mathbf{X}\mathbf{C})'(\mathbf{X}\mathbf{C}), \end{aligned}$$

and by Theorem 1.3, we have that

$$\begin{aligned}\text{rank}(\mathbf{B}'\mathbf{FB}) &= \text{rank}((\mathbf{XC})'(\mathbf{XC})) \\ &= \text{rank}(\mathbf{XC}) \\ &= s.\end{aligned}$$

### THEOREM 14.3

Set  $Q = (\mathbf{B}'\vec{\beta}_0 - \vec{m})'(\mathbf{B}'\mathbf{FB})^{-1}(\mathbf{B}'\vec{\beta}_0 - \vec{m})$ . Then, the following hold:

(1)  $Q/\sigma^2 \sim \chi^2(s, \lambda)$ , where

$$\lambda = \frac{1}{2\sigma^2}(\mathbf{B}'\vec{\beta} - \vec{m})'(\mathbf{B}'\mathbf{FB})^{-1}(\mathbf{B}'\vec{\beta} - \vec{m}).$$

(2)  $Q$  and SSE are independent.

**Proof:**

(1) The main idea is to use Theorem 4.1. Note that  $\vec{Y} \sim \mathcal{N}(\mathbf{X}\vec{\beta}, \sigma^2\mathbf{I})$ .

$$\begin{aligned}\mathbf{B}'\vec{\beta}_0 - \vec{m} &= \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}_0 - \vec{m} \\ &= \mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y} - \vec{m} \\ &\sim \text{MN}(\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta} - \vec{m}, \Sigma) \\ &= \text{MN}(\underbrace{\mathbf{B}'\vec{\beta} - \vec{m}}_{\vec{\mu}}, \Sigma),\end{aligned}$$

where

$$\begin{aligned}\Sigma &= \sigma^2\mathbf{B}'\mathbf{F}\mathbf{X}\mathbf{X}'(\mathbf{B}'\mathbf{F}\mathbf{X}')' \\ &= \sigma^2\mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{B}'\mathbf{F}'\mathbf{B} \\ &= \sigma^2\mathbf{B}'\mathbf{FB}.\end{aligned}$$

Therefore,

$$\mathbf{A}\Sigma = (\mathbf{B}'\mathbf{FB})^{-1}\sigma^2(\mathbf{B}'\mathbf{FB}) = \mathbf{I},$$

which is idempotent. It follows from Theorem 4.1 that  $Q \sim \chi^2(s, \lambda)$ , where  $\lambda$  is defined above.

(2)  $Q = (\mathbf{B}'\vec{\beta}_0 - \vec{m})'(\mathbf{B}'\mathbf{FB})^{-1}(\mathbf{B}'\vec{\beta}_0 - \vec{m})$ . Idea: rewrite  $Q$  into a quadratic form and use Theorem 5.1.

$$Q = (\mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y} - \vec{m})'(\mathbf{B}'\mathbf{FB})^{-1}(\mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y} - \vec{m})$$

Recall that SSE =  $\vec{Y}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}')\vec{Y}$ , and

$$\begin{aligned}\mathbf{B}'\vec{\beta}_0 &= \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\vec{\beta}_0 \\ &= \mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y} \\ &= (\mathbf{B}'\mathbf{F}\mathbf{X}')\vec{Y}.\end{aligned}$$

Hence,

$$\begin{aligned}(\mathbf{B}'\mathbf{F}\mathbf{X}')\sigma^2\mathbf{I}(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}') &= \sigma^2[\mathbf{B}'\mathbf{F}\mathbf{X}' - \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'] \\ &= \sigma^2[\mathbf{B}'\mathbf{F}\mathbf{X}' - \mathbf{B}'\mathbf{F}\mathbf{X}'] \\ &= \mathbf{O}.\end{aligned}$$

Therefore, by Theorem 5.1, we obtain the result.

Alternative Proof Idea: Prove that  $\mathbf{B}'\mathbf{F}\mathbf{X}'\vec{Y}$  is independent of SSE.

$$\begin{aligned}\mathbf{B}'\mathbf{F}\mathbf{X}'(\mathbf{I} - \mathbf{X}\mathbf{F}\mathbf{X}') &= \mathbf{B}'\mathbf{F}\mathbf{X}' - \mathbf{B}'\mathbf{F}\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}' \\ &= \mathbf{B}'\mathbf{F}\mathbf{X}.\end{aligned}$$

## General One-Way Classification Model

$$Y_{ij} = \mu + \alpha_i + \varepsilon_{ij},$$

where  $i = 1, \dots, a$  are the factors,  $j = 1, \dots, n_i$  are the levels, and  $n = n_1 + \dots + n_a$  are the total number of observations.

- $\mu$  is the global average;
- $\alpha_i$  is additional impact of group  $i$  ( $\alpha_i > 0$  higher than global average);
- $\varepsilon_{ij}$  is the error of group  $i$  on level  $j$ .

In matrix-vector form, we may write the model as follows.

$$\begin{pmatrix} Y_{11} \\ \vdots \\ Y_{1n_1} \\ \vdots \\ Y_{a1} \\ \vdots \\ Y_{an_a} \end{pmatrix} = \mathbf{X} \begin{pmatrix} \mu \\ \alpha_1 \\ \vdots \\ \alpha_a \end{pmatrix} + \vec{\varepsilon},$$

where

$$\mathbf{X} = \begin{pmatrix} n_1 \left\{ \begin{array}{cccc} 1 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 0 & \cdots & 0 \end{array} \right\} \\ n_2 \left\{ \begin{array}{cccc} 1 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 0 & \cdots & 0 \end{array} \right\} \\ \vdots \\ n_a \left\{ \begin{array}{cccc} 1 & 0 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & 1 \end{array} \right\} \end{pmatrix} \Rightarrow \mathbf{X}' = \begin{pmatrix} 1 & \cdots & 1 & 1 & \cdots & 1 & \cdots & 1 & \cdots & 1 \\ 1 & \cdots & 1 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1 & \cdots & 1 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 1 & \cdots & 1 \end{pmatrix}$$

and  $\text{rank}(\mathbf{X}) = a$ . Note that

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n & n_1 & n_2 & \cdots & n_a \\ n_1 & n_1 & 0 & \cdots & 0 \\ n_2 & 0 & n_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n_a & 0 & \cdots & 0 & n_a \end{pmatrix} \in \mathbb{R}^{(a+1) \times (a+1)}$$



A  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \begin{pmatrix} 0 & \cdots & 0 & 0 \\ \vdots & 1/n_1 & \ddots & \vdots \\ 0 & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1/n_a \end{pmatrix}$$

$$\mathbf{X}'\vec{Y} = \begin{pmatrix} \sum_{i=1}^a \sum_{j=1}^{n_i} Y_{ij} \\ \sum_{j=1}^{n_1} Y_{1j} \\ \vdots \\ \sum_{j=1}^{n_a} Y_{aj} \end{pmatrix} = \begin{pmatrix} Y_{..} \\ Y_{1.} \\ \vdots \\ Y_{a.} \end{pmatrix},$$

where  $Y_{..} = \sum_{i=1}^a \sum_{j=1}^{n_i} Y_{ij}$  and  $Y_{i.} = \sum_{j=1}^{n_i} Y_{ij}$ . Also,

$$\mathbf{F}\mathbf{X}' = \begin{pmatrix} 0 & \cdots & 0 & 0 \\ \vdots & 1/n_1 & \ddots & \vdots \\ 0 & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1/n_a \end{pmatrix} \begin{pmatrix} 1 & \cdots & 1 & 1 & \cdots & 1 & \cdots & 1 & \cdots & 1 \\ 1 & \cdots & 1 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1 & \cdots & 1 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 1 & \cdots & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 1/n_1 & \cdots & 1/n_1 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1/n_2 & \cdots & 1/n_2 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 1/n_a & \cdots & 1/n_a \end{pmatrix}$$

Therefore, a solution to the normal equation

$$\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}$$

is

$$\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y} = \begin{pmatrix} 0 \\ \frac{1}{n_1} Y_{1.} \\ \vdots \\ \frac{1}{n_a} Y_{a.} \end{pmatrix} = \begin{pmatrix} 0 \\ \bar{Y}_{1.} \\ \vdots \\ \bar{Y}_{a.} \end{pmatrix},$$

where  $\bar{Y}_{i.} = \frac{1}{n_i} Y_{i.}$ .

- Sum of Squares Total:

$$\text{SST} = \sum_{i=1}^a \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2$$

$$= \sum_{i=1}^a \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{n}.$$

- Sum of Squares Regression: Define  $\hat{Y} = \mathbf{X}\vec{\beta}_0 = \mathbf{X}\mathbf{F}\mathbf{X}'\vec{Y}$ . Hence,

$$\text{SSR} = \sum_{i=1}^a \sum_{j=1}^{n_i} (\hat{Y}_{ij} - \bar{Y}_{..})^2$$

$$= \hat{Y}'\hat{Y} - 2 \left( \sum_{i=1}^a \sum_{j=1}^{n_i} \hat{Y}_{ij} \right) \bar{Y}_{..} + n\bar{Y}_{..}^2.$$

Furthermore,

$$\begin{aligned}
 \sum_{i=1}^a \sum_{j=1}^{n_i} \hat{Y}_{ij} &= \vec{j}' \hat{\vec{Y}} \\
 &= \vec{j}' \mathbf{X} \vec{\beta}_0 \\
 &= (n \quad n_1 \quad \cdots \quad n_a) \vec{\beta}_0 \\
 &= n_1 \bar{Y}_{1.} + n_2 \bar{Y}_{2.} + \cdots + n_a \bar{Y}_{a.} \\
 &= Y_{..} \\
 &= n \bar{Y}_{..}
 \end{aligned}$$

Furthermore,  $SSR = \hat{\vec{Y}}' \hat{\vec{Y}} - n \bar{Y}_{..}^2$ , and

$$\hat{\vec{Y}}' = \mathbf{X} \vec{\beta}_0 = \mathbf{X} \begin{pmatrix} 0 \\ \bar{Y}_{1.} \\ \vdots \\ \bar{Y}_{a.} \end{pmatrix} = \begin{pmatrix} n_1 \begin{Bmatrix} \bar{Y}_{1.} \\ \vdots \\ \bar{Y}_{1.} \end{Bmatrix} \\ \vdots \\ n_a \begin{Bmatrix} \bar{Y}_{a.} \\ \vdots \\ \bar{Y}_{a.} \end{Bmatrix} \end{pmatrix}$$

Therefore,

$$\begin{aligned}
 SSR &= n_1 \bar{Y}_{1.}^2 + \cdots + n_a \bar{Y}_{a.}^2 - n \bar{Y}_{..}^2 \\
 &= \sum_{i=1}^a \frac{Y_{i.}^2}{n_i} - \frac{Y_{..}^2}{n}.
 \end{aligned}$$

- Sum of Squares Error:

$$\begin{aligned}
 SSE &= SST - SSR \\
 &= \sum_{i=1}^a \sum_{j=1}^{n_i} Y_{ij}^2 - \sum_{i=1}^a \frac{Y_{i.}^2}{n_i}.
 \end{aligned}$$

By Theorem 13.1 and Theorem 13.2, we have the following distributions.

- $SSE/\sigma^2 \sim \chi^2(n - a)$ .
- $SSR/\sigma^2 \sim \chi^2(a - 1, \lambda)$ , where

$$\lambda = \frac{1}{2\sigma^2} (\mathbf{X} \vec{\beta})' (\mathbf{X} \mathbf{F} \mathbf{X}' - \frac{1}{n} \mathbf{J}) \mathbf{X} \vec{\beta}.$$

LECTURE 15  
6th March

## 15 Lecture 15: ANOVA Table For One-Way Classification

For  $H_0: \mathbf{X} \vec{\beta} = \vec{0}$  versus  $H_A: \mathbf{X} \vec{\beta} \neq \vec{0}$ .

| Source of Variation | Degrees of Freedom | Sum of Squares | Mean Square | $F$     |
|---------------------|--------------------|----------------|-------------|---------|
| Due to Regression   | $a - 1$            | SSR            | MSR         | MSR/MSE |
| Error               | $n - a$            | SSE            | MSE         |         |
| Total               | $n - 1$            | SST            |             |         |

if  $F = MSR/MSE > F_\alpha(a - 1, n - a)$ , then we conclude that the model  $Y_{ij} + \mu + \alpha_i + \varepsilon_{ij}$  accounts for significantly more variation in the  $Y$  variable than the model  $Y_{ij} + \mu + \varepsilon_{ij}$ .

## Estimable Linear Combinations

$\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}'$ , where  $\vec{b} \in \mathbb{R}^{a+1}$ .

$$\begin{aligned} \mathbf{F}\mathbf{X}'\mathbf{X} &= \begin{pmatrix} 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 1/n_1 & \cdots & 1/n_1 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1/n_2 & \cdots & 1/n_2 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & 1/n_a & \cdots & 1/n_a \end{pmatrix} \\ &\times \begin{pmatrix} 1 & \cdots & 1 & 1 & \cdots & 1 & \cdots & 1 & \cdots & 1 \\ 1 & \cdots & 1 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1 & \cdots & 1 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 1 & \cdots & 1 \end{pmatrix} \mathbf{X} \\ &= \text{exercise} \\ &= \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ 1 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & 0 \cdots & 1 \end{pmatrix}. \end{aligned}$$

$$\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = (b_1 + \cdots + b_a \quad b_1 \quad b_2 \quad \cdots \quad b_a) = (b_0 \quad b_1 \quad \cdots \quad b_a),$$

where  $b_0 = b_1 + \cdots + b_a$ . Now,

$$\begin{aligned} \vec{b}'\vec{\beta} &= (b_0 \quad \cdots \quad b_a) \begin{pmatrix} \mu \\ \alpha_1 \\ \vdots \\ \alpha_a \end{pmatrix} \\ &= b_0\vec{\mu} + b_1\alpha_1 + \cdots + b_a\alpha_a \\ &= (b_1 + \cdots + b_a)\vec{\mu} + b_1\alpha_1 + \cdots + b_a\alpha_a \\ &= b_1(\mu + \alpha_1) + \cdots + b_a(\mu + \alpha_a). \end{aligned}$$

Questions:

- (1) Is  $\mu$  estimable? No.
- (2) Are  $\alpha_i$ 's estimable? No.  $b_i = 1, b_j = 0$  for  $i \neq j$ , so we will get  $\mu + \alpha_i \neq \alpha_i$ .
- (3) Is  $\mu + \alpha_i$  estimable? Yes. Choose  $b_i = 1$  and  $b_j = 0$  for  $i \neq j$ .
- (4) Is  $\alpha_i - \alpha_j$  estimable (for  $i \neq j$ )? Yes.  $b_i = 1, b_j = -1$ , and  $b_k = 0$  for  $k \neq i, j$ .

## Confidence Interval

If  $\vec{b}'\vec{\beta}$  is estimable, then a  $100(1 - \alpha)\%$  confidence interval for  $\vec{b}'\vec{\beta}$  is

$$\vec{b}'\vec{\beta}_0 \pm t_{n-a, \alpha/2} \hat{\sigma} \sqrt{\vec{b}'\mathbf{F}\vec{b}}.$$

To get a CI for  $\beta_i - \beta_j$ , we need to consider

$$\vec{b}' = \left( 0 \quad \cdots \quad 0 \quad \underbrace{1}_i \quad \cdots \quad \underbrace{-1}_j \quad 0 \quad \cdots \quad 0 \right),$$

we have  $\vec{b}'\vec{\beta} = \vec{\beta}_i - \vec{\beta}_j$ . Hence,

$$\begin{aligned}\vec{b}'\mathbf{F}\vec{b} &= (0 \quad \cdots \quad 0 \quad 1/n_i \quad \cdots \quad -1/n_j \quad 0 \quad \cdots \quad 0) \vec{b} \\ &= \frac{1}{n_i} + \frac{1}{n_j}.\end{aligned}$$

Therefore,  $\hat{\sigma}\sqrt{\frac{1}{n_i} + \frac{1}{n_j}}$  is the standard error for the CI of  $\beta_i - \beta_j$ , and a  $100(1 - \alpha)\%$  confidence interval for  $\beta_i - \beta_j$  is

$$\beta_i^0 - \beta_j^0 \pm t_{n-a,\alpha/2}\hat{\sigma}\sqrt{\frac{1}{n_i} + \frac{1}{n_j}}.$$

## Two-way Nested Classification Part I

$$Y_{ijk} = \mu + \alpha_i + \beta_{ij} + \varepsilon_{ijk},$$

where  $i = 1, \dots, a$  (category  $i$ ),  $j = 1, \dots, b_i$  (subclass  $j$ ),  $k = 1, \dots, n_{ij}$  (observation  $k$ ).

- $\mu$  is the overall mean;
- $\alpha_i$  is the effect due to category  $i$ ;
- $\beta_{ij}$  is the effect due to subclass  $j$ ;
- $\varepsilon_{ijk}$  is the error.

In matrix-vector form, we may write the model as follows (why would anyone write this).

$$\underbrace{\begin{pmatrix} Y_{111} \\ \vdots \\ Y_{11n_{11}} \\ \vdots \\ Y_{1b_11} \\ \vdots \\ Y_{1b_1n_{1b_1}} \\ \vdots \\ Y_{211} \\ \vdots \\ Y_{21n_{21}} \\ \vdots \\ Y_{2b_21} \\ \vdots \\ Y_{2b_2n_{2b_2}} \\ \vdots \\ Y_{a11} \\ \vdots \\ Y_{a1n_{a1}} \\ \vdots \\ Y_{ab_a1} \\ \vdots \\ Y_{ab_an_{ab_a}} \end{pmatrix}}_{\vec{Y}} = \mathbf{X} \underbrace{\begin{pmatrix} \mu \\ \alpha_1 \\ \vdots \\ \alpha_a \\ \beta_{11} \\ \vdots \\ \beta_{1b_1} \\ \vdots \\ \beta_{a1} \\ \vdots \\ \beta_{ab_a} \end{pmatrix}}_{\vec{\beta}} + \vec{\varepsilon},$$

where I personally can't be bothered to write out  $\mathbf{X}$ .

Notes:

- Total number of observations:  $n = \sum_{i=1}^a \sum_{j=1}^{b_i} n_{ij} = \sum_{i=1}^a n_{i.}$
- Total number of parameters:  $m = 1 + a + \sum_{i=1}^a b_i.$
- $\text{rank}(\mathbf{X}'\mathbf{X}) = b.$

### EXAMPLE 15.1

Suppose we want to know the students opinion of the instructor's classroom use of computer facility (0  $\leftrightarrow$  10). For example, if we have two courses English (two sections) and Geology (three sections):

- English Section 1: 5.
- English Section 2: 8, 10, 9.
- Geology Section 1: 8, 10.
- Geology Section 2: 6, 2, 1, 3.
- Geology Section 3: 3, 7.

We calculate the following.

- $a = 2, b_1 = 2, b_2 = 3.$
- $n_{11} = 1, n_{12} = 3, n_{21} = 2, n_{22} = 4, n_{23} = 2.$
- $n = n_{11} + n_{12} + n_{21} + n_{22} + n_{23} = 1 + 3 + 2 + 4 + 2 = 12.$
- $m = 1 + a + b_1 + b_2 = 1 + 2 + 2 + 3 = 8.$

$$\vec{\beta} = \begin{pmatrix} \mu \\ \alpha_1 \\ \alpha_2 \\ \beta_{11} \\ \beta_{12} \\ \beta_{21} \\ \beta_{22} \\ \beta_{23} \end{pmatrix}$$

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n_{..} & n_{1.} & n_{2.} & n_{11} & n_{12} & n_{21} & n_{22} & n_{23} \\ n_{1.} & n_{2.} & 0 & n_{11} & n_{12} & 0 & 0 & 0 \\ n_{2.} & 0 & n_{2.} & 0 & 0 & n_{21} & n_{22} & n_{23} \\ n_{11} & n_{11} & 0 & n_{11} & 0 & 0 & 0 & 0 \\ n_{12} & n_{12} & 0 & 0 & n_{12} & 0 & 0 & 0 \\ n_{21} & 0 & n_{21} & 0 & 0 & n_{21} & 0 & 0 \\ n_{22} & 0 & n_{22} & 0 & 0 & 0 & n_{22} & 0 \\ n_{23} & 0 & n_{23} & 0 & 0 & 0 & 0 & n_{23} \end{pmatrix}.$$

$$\mathbf{X}'\vec{Y} = (Y_{..} \quad Y_{1.} \quad Y_{2.} \quad Y_{11} \quad Y_{12} \quad Y_{21} \quad Y_{22} \quad Y_{23}).$$

$\text{rank}(\mathbf{X}'\mathbf{X}) = b_1 + b_2 = 5$ , and a  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \begin{pmatrix} \mathbf{O} & \mathbf{O} \\ \mathbf{O} & \text{diag}(1/n_{11}, \dots, /n_{23}) \end{pmatrix}.$$

$$\mathbf{X} = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Therefore,

$$\vec{\beta}'_0 = \mathbf{F}\mathbf{X}'\vec{Y} = (0 \quad 0 \quad 0 \quad \bar{Y}_{11} \quad \bar{Y}_{12} \quad \bar{Y}_{21} \quad \bar{Y}_{22} \quad \bar{Y}_{23})'$$

LECTURE 16  
9th March

## 16 Lecture 16: Two-Way Nested Classification Part II

- Model 1:  $Y_{ijk} = \mu + \varepsilon_{ijk}$ .
- Model 2:  $Y_{ijk} = \mu + \alpha_i + \varepsilon_{ijk}$ .
- Model 3:  $Y_{ijk} = \mu + \alpha_i + \beta_{ij} + \varepsilon_{ijk}$ .

### ANOVA

$$\begin{aligned} \text{SST} &= \sum_{i=1}^a \sum_{j=1}^{b_i} \sum_{k=1}^{n_{ij}} (Y_{ijk} - \bar{Y}_{...})^2 \\ &= \vec{Y}'\vec{Y} - n\bar{Y}_{...}^2 \\ \text{SSR} &= \sum_{i=1}^a \sum_{j=1}^{b_i} \sum_{k=1}^{n_{ij}} (\hat{Y}_{ijk} - \bar{Y}_{...})^2 \\ &= \vec{\beta}'_0 \mathbf{X}'\vec{Y} - n\bar{Y}_{...}^2 \\ &= \text{SS}(\alpha, \beta : \alpha | \mu) \\ \text{SSE} &= \text{SST} - \text{SSR} \\ &= \sum_{i=1}^a \sum_{j=1}^{b_i} \left( \sum_{k=1}^{n_{ij}} Y_{ijk}^2 - \frac{1}{n_{ij}} Y_{ij.}^2 \right) \end{aligned}$$

### ANOVA Table I

Testing the overall effectiveness of Model 3.

| Source of Variation                          | Degrees of Freedom | Sum of Squares | Mean Square | $F$     |
|--|--------------------|----------------|-------------|---------|
| Fitting $\alpha, \beta : \alpha$ after $\mu$ | $b - 1$            | SSR            | MSR         | MSR/MSE |
| Error  | $n - b$            | SSE            | MSE         |         |
| Total  | $n - 1$            | SST            |             |         |

Tests on the significant impact on the variation in  $Y$ .

## ANOVA Table II

| Source of Variation                          | Degrees of Freedom | Sum of Squares                     | Mean Square                         | $F$                               |
|--|--------------------|------------------------------------|-------------------------------------|-----------------------------------|
| Fitting $\alpha$ after $\mu$                 | $a - 1$            | $SS(\alpha   \mu)$                 | $MSR(\alpha   \mu)$                 | $F(\alpha   \mu)$                 |
| Fitting $\beta : \alpha$ after $\mu, \alpha$ | $b - a$            | $SS(\beta : \alpha   \mu, \alpha)$ | $MSR(\beta : \alpha   \mu, \alpha)$ | $F(\beta : \alpha   \mu, \alpha)$ |
| Error  | $n - b$            | SSE                                | MSE                                 |                                   |
| Total  | $n - 1$            | SST                                |                                     |                                   |

- $SS(\alpha | \mu) = SSR$  of Model 2; that is, we are checking the impact of  $\alpha$  on the variance.
- $SS(\beta : \alpha | \mu, \alpha) = SS(\alpha, \beta : \alpha | \mu) - SS(\alpha | \mu)$ ; that is, we are checking the impact of  $\beta$  after  $\alpha$  on the model.

$$F(\alpha | \mu) = \frac{MSR(\alpha | \mu)}{MSE} \sim F(\alpha - 1, n - b).$$

$$F(\beta : \alpha | \mu, \alpha) = \frac{MSR(\beta : \alpha)}{MSE} \sim F(b - a, n - b).$$

### EXAMPLE 16.1

Refer to the English and Geology data from Example 15.1.

- $SST = 110$
- $SSR = 84$ .
- $SSE = 26$ .
- $SS(\alpha | \mu) = \sum_{i=1}^a \frac{Y_{i..}^2}{n_i} - \frac{Y_{...}^2}{n} = 24$ .
- $SS(\beta : \alpha | \mu, \alpha) = 84 - 24 = 60$ .

ANOVA Table I:

| Source of Variation                          | Degrees of Freedom | Sum of Squares | Mean Square | $F$               |
|--|--------------------|----------------|-------------|-------------------|
| Fitting $\alpha, \beta : \alpha$ after $\mu$ | 4                  | 84             | 21          | $21(7)/26 = 5.66$ |
| Error  | 7                  | 26             | $26/7$      |                   |
| Total  | 11                 | 110            |             |                   |

ANOVA Table II:

| Source of Variation                          | Degrees of Freedom | Sum of Squares | Mean Square | $F$ |
|--|--------------------|----------------|-------------|-----|
| Fitting $\alpha$ after $\mu$                 | 1                  | 24             | 24          | 6.5 |
| Fitting $\beta : \alpha$ after $\mu, \alpha$ | 3                  | 60             | 20          | 5.4 |
| Error  | 7                  | 26             | $26/7$      |     |
| Total  | 11                 | 110            |             |     |

- Testing  $\alpha, \beta : \alpha$  after  $\mu$ :  $5.66 > F_{0.05}(4, 7) = 4.12$ , reject the fact that we do not need  $\alpha$  and  $\beta$ , so Model 3 is adequate. Model 3 accounts for significantly more variation than Model 1.
- Testing  $\alpha$  after  $\mu$ :  $6.5 > F_{0.05}(1, 7) = 5.59$ , we need  $\alpha$ . Model 2 accounts for significantly more variation than Model 1.
- Testing  $\beta : \alpha$  after  $\mu, \alpha$ :  $5.4 > F_{0.05}(3, 7) = 4.35$ , we need  $\beta$ . Model 3 accounts for significantly more variation than Model 2.

## Estimable Combinations

$\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}'$ . Write  $\vec{b}'\mathbf{F}\mathbf{X}' = \vec{c}'$ . Also,  $\vec{b}'\vec{\beta} = \vec{c}'\mathbf{X}\vec{\beta}$ . Idea:

$$\begin{aligned}\vec{c}' &= (c_{111} \quad \cdots \quad c_{11n_{11}} \quad \cdots \quad c_{ij1} \quad \cdots \quad c_{ijn_{ij}} \quad \cdots). \\ \vec{c}'\mathbf{X} &= (w_{11} \quad \cdots \quad w_{1b_1} \quad w_{21} \quad \cdots \quad w_{2b_2} \quad \cdots \quad w_{a1} \quad \cdots \quad w_{ab_a}). \\ \vec{b}'\vec{\beta} &= (w_{11} \quad \cdots \quad w_{1b_1} \quad \cdots) \vec{\beta} \\ &= \sum_{i=1}^a \sum_{j=1}^{b_i} w_{ij} (\mu + \alpha_i + \beta_{ij}).\end{aligned}$$

Conclusion:

- (1)  $\mu, \mu + \alpha_i, \alpha_i$  are not estimable.
- (2)  $\mu + \alpha_i + \beta_{ij}$  is estimable.
- (3)  $\beta_{ij} - \beta_{i\ell}$  is estimable.
- (4)  $\mu + \alpha_i + \sum_{j=1}^{b_i} w_{ij} \beta_{ij}$  is estimable if  $\sum_{j=1}^{b_i} w_{ij} = 1$ .
- (5)  $\alpha_i - \alpha_\ell + \sum_{j=1}^{b_i} w_{ij} \beta_{ij} - \sum_{\ell=1}^{b_\ell} w_{i\ell} \beta_{i\ell}$  is estimable if  $\sum_{j=1}^{b_i} w_{ij} = \sum_{\ell=1}^{b_\ell} w_{i\ell} = 1$ .
- (6)  $\alpha_i - \alpha_\ell$  is not estimable.

## Hypothesis Testing

- $H_0: \beta_{i1} = \cdots = \beta_{ib_i}, i = 1, \dots, a;$
- $H_A: \text{at least one equality fails.}$

Define  $(b_1 - 1) + (b_2 - 1) + \cdots + (b_a - 1) = n'$  and  $m = 1 + a + \sum_{i=1}^a b_i$ . Hence,  $\mathbf{B}' \in \mathbb{R}^{n' \times m}$ , where  $\text{rank}(\mathbf{B}) = b - a$ . Therefore,

$$\mathbf{B}' = (\mathbf{O}_{n' \times (a+1)} \quad \text{diag}(\mathbf{Z}_{b_1}, \dots, \mathbf{Z}_{b_a})),$$

where  $\mathbf{Z}_{b_i} = \begin{pmatrix} \mathbf{1}_{(b_i-1) \times 1} & \text{diag}(\underbrace{-1, \dots, -1}_{b_i-1}) \end{pmatrix} \in \mathbb{R}^{(b_i-1) \times b_i}$  for  $i = 1, \dots, a$ .

$$Q = (\mathbf{B}'\vec{\beta}_0)'(\mathbf{B}'\mathbf{F}\mathbf{B})(\mathbf{B}'\vec{\beta}_0) = \text{SS}(\beta : \alpha \mid \mu, \alpha).$$

$$F = \frac{Q/(n-a)}{\text{SSE}/(n-b)} \sim F(b-a, n-b).$$

### REMARK

If  $n_{ij} = r$  for all  $i, j$ , and  $b_i = c$  for all  $i$ , then the model is called a **balanced model**.

From Example 15.1, we have

$$\vec{\beta} = \begin{pmatrix} \mu \\ \alpha_1 \\ \alpha_2 \\ \beta_{11} \\ \beta_{12} \\ \beta_{21} \\ \beta_{22} \\ \beta_{23} \end{pmatrix},$$



$$\vec{b}' = (a_1 \quad a_2 \quad a_3 \quad b_{11} \quad b_{12} \quad b_{21} \quad b_{22} \quad b_{23}).$$

Therefore,  $\vec{b}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{b}'$ , where

$$a_1 = b_{11} + b_{12} + b_{21} + b_{22} + b_{23},$$

$$a_2 = b_{11} + b_{12},$$

$$a_3 = b_{21} + b_{22} + b_{23}.$$

Therefore,

$$\begin{aligned} \vec{b}'\vec{\beta} &= a_1\mu + a_2\alpha_1 + a_3\alpha_2 + b_{11}\beta_{11} + b_{12}\beta_{12} + b_{21}\beta_{21} + b_{22}\beta_{22} + \beta_{23}\beta_{23} \\ &= b_{11}(\mu + \alpha_1 + \beta_{11}) + b_{12}(\mu + \alpha_1 + \beta_{12}) + b_{21}(\mu + \alpha_2 + \beta_{21}) + b_{22}(\mu + \alpha_2 + \beta_{22}) + b_{23}(\mu + \alpha_2 + \beta_{23}). \end{aligned}$$

LECTURE 17  
13th March

## 17 Lecture 17: Two-way Crossed Classification (No Interaction) Part I

Basic Data Structure:

- Two factors  $A$  and  $B$ , where factor  $A$  has  $a$  levels, and factor  $B$  has  $b$  levels.

| $A \setminus B$ | 1        | 2        | ...      | $b$      |
|-----------------|----------|----------|----------|----------|
| 1               | $n_{11}$ | $n_{12}$ | ...      | $n_{1b}$ |
| 2               | $n_{21}$ | $n_{22}$ | ...      | $n_{2b}$ |
| $\vdots$        | $\vdots$ | $\vdots$ | $\ddots$ | $\vdots$ |
| $a$             | $n_{a1}$ | $n_{a2}$ | ...      | $n_{ab}$ |

Let  $n_{ij}$  be the number of observations at cell  $(i, j)$ . The model with **no interaction** is defined as

$$Y_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}.$$

Write

$$\vec{\beta} = (\mu \quad \alpha_1 \quad \cdots \quad \alpha_a \quad \beta_1 \quad \cdots \quad \beta_b)'$$

For this model,  $n_{ij} = 0$  or  $1$ . If  $n_{ij} = 1$  for all cells, the model is said to be **balanced**. In this case, we have

$$\mathbf{X} = \begin{matrix} (i,j) \\ (1,1) \\ \vdots \\ (1,b) \\ (2,1) \\ \vdots \\ (2,b) \\ \vdots \\ (a,1) \\ \vdots \\ (a,b) \end{matrix} \begin{pmatrix} \mu & \alpha_1 & \alpha_2 & \cdots & \alpha_a & \beta_1 & \beta_2 & \cdots & \beta_b \\ 1 & 1 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 1 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 1 & \cdots & 0 & 0 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 1 & \cdots & 1 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 1 & \cdots & 1 & 0 & 0 & \cdots & 1 \end{pmatrix} \in \mathbb{R}^{(ab) \times \overbrace{(1+a+b)}^m}.$$

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n_{..} & n_{1.} & n_{2.} & \cdots & n_{a.} & n_{.1} & \cdots & n_{.b} \\ n_{1.} & n_{11} & 0 & \cdots & 0 & 1 & \cdots & 1 \\ n_{2.} & 0 & n_{22} & \ddots & 0 & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ n_{a.} & 0 & 0 & \cdots & n_{a.} & 1 & \cdots & 1 \\ n_{.1} & 1 & 1 & \cdots & 1 & n_{.1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ n_{.b} & 1 & 1 & \cdots & 1 & 0 & \cdots & n_{.b} \end{pmatrix} \in \mathbb{R}^{m \times m}.$$

For a general model  $n_{ij} = 0$  or 1, we have

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n_{..} & n_{1.} & n_{2.} & \cdots & n_{a.} & n_{.1} & \cdots & n_{.b} \\ n_{1.} & n_{11} & 0 & \cdots & 0 & n_{11} & \cdots & n_{1b} \\ n_{2.} & 0 & n_{22} & \ddots & 0 & n_{21} & \cdots & n_{2b} \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots \\ n_{a.} & 0 & 0 & \cdots & n_{aa} & n_{a1} & \cdots & n_{ab} \\ n_{.1} & n_{11} & n_{21} & \cdots & n_{a1} & n_{.1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \ddots & \vdots \\ n_{.b} & n_{1b} & n_{2b} & \cdots & n_{ab} & 0 & \cdots & n_{.b} \end{pmatrix}.$$

$$\mathbf{X}'\vec{Y} = \begin{pmatrix} Y_{..} \\ Y_{1.} \\ \vdots \\ Y_{a.} \\ Y_{.1} \\ \vdots \\ Y_{.b} \end{pmatrix}.$$

The normal equation is  $\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}$ . We note that  $\text{rank}(\mathbf{X}) = a + b - 1$  since the sum of rows 2 to  $a$  equals the first row, and the sum of rows  $a + 2$  to  $a + b + 1$  also equals to the first row. Since the number of parameters is  $a + b + 1$ , it follows that we have the freedom of removing two equations.

General Rules:

- (1) Remove the first equation by setting  $\mu = 0$ .
- (2) If  $a < b$ , set  $\alpha_1 = 0$ . If  $a > b$ , set  $\beta_b = 0$ . If  $a = b$ , set either  $\alpha_1 = 0$  or  $\beta_b = 0$ .

Let

$$\begin{aligned} \mathbf{D}_{a.} &= \text{diag}(n_{11}, \dots, n_{aa}), \\ \mathbf{D}_{.b} &= \text{diag}(n_{.1}, \dots, n_{.b}), \\ \mathbf{D}_{.(b-1)} &= \text{diag}(n_{.1}, \dots, n_{.(b-1)}), \\ \mathbf{N}_{ab} &= \begin{pmatrix} n_{11} & \cdots & n_{1b} \\ \vdots & \ddots & \vdots \\ n_{a1} & \cdots & n_{ab} \end{pmatrix}. \end{aligned}$$

Therefore,

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n_{..} & n_{1.} & \cdots & n_{a.} & n_{.1} & \cdots & n_{.b} \\ n_{1.} & & & & & & \\ \vdots & & \mathbf{D}_{a.} & & & & \mathbf{N}_{ab} \\ n_{a.} & & & & & & \\ n_{.1} & & & & & & \\ \vdots & & \mathbf{N}'_{ab} & & & & \mathbf{D}_{.b} \\ n_{.b} & & & & & & \end{pmatrix}.$$

Furthermore,

$$\vec{Y}_{a.} = (Y_{1.} \quad \cdots \quad Y_{a.})', \quad \vec{Y}_{.(b-1)} = (Y_{.1} \quad \cdots \quad Y_{.(b-1)})'.$$

Removing  $\mu$  and  $\beta_b$  (first row and last row are linear combination of other rows), we obtain

$$\begin{pmatrix} \mathbf{D}_{a.} & \mathbf{N}_{a(b-1)} \\ \mathbf{N}'_{a(b-1)} & \mathbf{D}_{.(b-1)} \end{pmatrix} \begin{pmatrix} \vec{\alpha} \\ \vec{\beta}_{b-1} \end{pmatrix} = \begin{pmatrix} \vec{Y}_{a.} \\ \vec{Y}_{.(b-1)} \end{pmatrix},$$

where  $\vec{\alpha} = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_a \end{pmatrix}$ , and  $\vec{\beta}_{b-1} = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_{b-1} \end{pmatrix}$ . Set  $\mathbf{N} = \mathbf{N}_{a(b-1)}$ . Therefore,

$$\begin{aligned} \mathbf{D}_a \vec{\alpha} + \mathbf{N} \vec{\beta}_{b-1} &= \vec{Y}_a \\ \mathbf{N}' \vec{\alpha} + \mathbf{D}_{(b-1)} \vec{\beta}_{b-1} &= \vec{Y}_{(b-1)} \end{aligned}$$

From the first equation, we obtain

$$\vec{\alpha} = \mathbf{D}_a^{-1} (\vec{Y}_a - \mathbf{N} \vec{\beta}_{b-1}).$$

Substituting  $\vec{\alpha}$  into the second equation, we get

$$\begin{aligned} \mathbf{N}' \mathbf{D}_a^{-1} (\vec{Y}_a - \mathbf{N} \vec{\beta}_{b-1}) + \mathbf{D}_{(b-1)} \vec{\beta}_{b-1} &= \vec{Y}_{(b-1)} \\ \mathbf{N}' \mathbf{D}_a^{-1} \vec{Y}_a - \mathbf{N}' \mathbf{D}_a^{-1} \mathbf{N} \vec{\beta}_{b-1} + \mathbf{D}_{(b-1)} \vec{\beta}_{b-1} &= \vec{Y}_{(b-1)} \\ -\mathbf{N}' \mathbf{D}_a^{-1} \mathbf{N} \vec{\beta}_{b-1} + \mathbf{D}_{(b-1)} \vec{\beta}_{b-1} &= \vec{Y}_{(b-1)} - \mathbf{N}' \mathbf{D}_a^{-1} \vec{Y}_a \\ (\mathbf{D}_{(b-1)} - \mathbf{N}' \mathbf{D}_a^{-1} \mathbf{N}) \vec{\beta}_{b-1} &= \vec{Y}_{(b-1)} - \mathbf{N}' \mathbf{D}_a^{-1} \vec{Y}_a. \end{aligned}$$

Define

$$\begin{aligned} \mathbf{C} &= \mathbf{D}_{(b-1)} - \mathbf{N}' \mathbf{D}_a^{-1} \mathbf{N}, \\ \vec{r} &= \vec{Y}_{(b-1)} - \mathbf{N}' \mathbf{D}_a^{-1} \vec{Y}_a, \\ \mathbf{M} &= \mathbf{D}_a^{-1} \mathbf{N}, \end{aligned}$$

so solving for  $\vec{\beta}_{b-1}$  (assuming that  $\mathbf{C}^{-1}$  exists) above yields

$$\vec{\beta}_{b-1} = \mathbf{C}^{-1} \vec{r}.$$

Furthermore,

$$\vec{\alpha} = \mathbf{D}_a^{-1} \vec{Y}_a - \mathbf{M} \mathbf{C}^{-1} \vec{r}.$$

Hence, one solution is

$$\vec{\beta}_0 = \begin{pmatrix} 0 \\ \vec{Y}_a - \mathbf{M} \mathbf{C}^{-1} \vec{r} \\ \mathbf{C}^{-1} \vec{r} \\ 0 \end{pmatrix} \begin{matrix} \mu \\ \vec{\alpha} \\ \vec{\beta}_{b-1} \\ \beta_b \end{matrix},$$

where  $\mathbf{D}_a^{-1} \vec{Y}_a = \vec{Y}_a$ .

A  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \begin{pmatrix} 0 & \mathbf{O} & \mathbf{O} & 0 \\ \mathbf{O} & \mathbf{D}_a^{-1} + \mathbf{M} \mathbf{C}^{-1} \mathbf{M}' & -\mathbf{M} \mathbf{C}^{-1} & \mathbf{O} \\ \mathbf{O} & -\mathbf{C}^{-1} \mathbf{M}' & \mathbf{C}^{-1} & \mathbf{O} \\ 0 & \mathbf{O} & \mathbf{O} & 0 \end{pmatrix}.$$

To verify this, we must check the following (exercise).

$$\begin{aligned} & \begin{pmatrix} \mathbf{D}_a^{-1} + \mathbf{M} \mathbf{C}^{-1} \mathbf{M}' & -\mathbf{M} \mathbf{C}^{-1} \\ -\mathbf{C}^{-1} \mathbf{M}' & \mathbf{C}^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{D}_a & \mathbf{N} \\ \mathbf{N}' & \mathbf{D}_{(b-1)} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{I}_a + \mathbf{M} \mathbf{C}^{-1} \mathbf{M}' \mathbf{D}_a & -\mathbf{M} \mathbf{C}^{-1} \mathbf{N}' & (\mathbf{D}_a^{-1} + \mathbf{M} \mathbf{C}^{-1} \mathbf{M}') \mathbf{N} - \mathbf{M} \mathbf{C}^{-1} \mathbf{D}_{(b-1)} \\ -\mathbf{C}^{-1} \mathbf{M}' \mathbf{D}_a & + \mathbf{C}^{-1} \mathbf{N}' & -\mathbf{C}^{-1} \mathbf{M}' \mathbf{N} + \mathbf{C}^{-1} \mathbf{D}_{(b-1)} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{I}_a & \mathbf{O} \\ \mathbf{O} & \mathbf{I}_{b-1} \end{pmatrix} \\ &= \mathbf{I}_{a+b-1}. \end{aligned}$$

## 18 Lecture 18: Two-way Crossed Classification (No Interaction) Part II

We know that  $\sum_{i=1}^a \alpha_i = 0$  and  $\sum_{j=1}^b \beta_j = 0$ .

- $$\text{SST} = \sum_{i=1}^a \sum_{j=1}^b Y_{ij}^2 - \frac{Y_{..}^2}{n}.$$

$$\begin{aligned} \text{SS}(\alpha, \beta \mid \mu) &= \vec{\beta}'_0 \mathbf{X}' \vec{Y} - \frac{Y_{..}^2}{n} \\ &= \vec{\beta}'_0 \begin{pmatrix} Y_{..} \\ \vec{Y}_{a.} \\ \vec{Y}_{.(b-1)} \\ Y_{.b} \end{pmatrix} - \frac{Y_{..}^2}{n} \\ &= \begin{pmatrix} 0 & (\vec{Y}_{a.} - \mathbf{M}\mathbf{C}^{-1}\vec{r})' & (\mathbf{C}^{-1}\vec{r})' & 0 \end{pmatrix} \begin{pmatrix} Y_{..} \\ \vec{Y}_{a.} \\ \vec{Y}_{.(b-1)} \\ Y_{.b} \end{pmatrix} - \frac{Y_{..}^2}{n} \\ &= (\mathbf{D}_a^{-1}\vec{Y}_{a.} - \mathbf{M}\mathbf{C}^{-1}\vec{r})'\vec{Y}_{a.} + (\mathbf{C}^{-1}\vec{r})'\vec{Y}_{.(b-1)} - \frac{Y_{..}^2}{n} \\ &= \vec{Y}'_{a.} \mathbf{D}_a^{-1} \vec{Y}_{a.} - (\mathbf{M}\mathbf{C}^{-1}\vec{r})'\vec{Y}_{a.} + (\mathbf{C}^{-1}\vec{r})'\vec{Y}_{.(b-1)} - \frac{Y_{..}^2}{n} \\ &= \sum_{i=1}^a \frac{Y_{i.}^2}{n_{i.}} + (\mathbf{C}^{-1}\vec{r})'(\vec{Y}_{.(b-1)} - \mathbf{M}'\vec{Y}_{a.}) - \frac{Y_{..}^2}{n} \\ &= \sum_{i=1}^a \frac{Y_{i.}^2}{n_{i.}} + \vec{\beta}'_{b-1} \vec{r} - \frac{Y_{..}^2}{n}, \end{aligned}$$

where  $\mathbf{C}^{-1}\vec{r} = \vec{\beta}_{b-1}$ , and  $\vec{r} = \vec{Y}_{.(b-1)} - \mathbf{M}'\vec{Y}_{a.}$ .

### ANOVA Table I

| Source of Variation                 | Degrees of Freedom | Sum of Squares                      | Mean Square                         | F                           |
|-------------------------------------|--------------------|-------------------------------------|-------------------------------------|-----------------------------|
| Fitting $\alpha, \beta$ after $\mu$ | $a + b - 2$        | $\text{SS}(\alpha, \beta \mid \mu)$ | $\text{MS}(\alpha, \beta \mid \mu)$ | $F(\alpha, \beta \mid \mu)$ |
| Error                               | $n - (a + b - 1)$  | SSE                                 | MSE                                 |                             |
| Total                               | $n - 1$            | SST                                 |                                     |                             |

- $$\text{SS}(\alpha, \beta \mid \mu) = \sum_{i=1}^a \frac{Y_{i.}^2}{n_{i.}} + \vec{\beta}'_{b-1} \vec{r} - \frac{Y_{..}^2}{n}.$$

### ANOVA Table II

| Source of Variation                 | Degrees of Freedom | Sum of Squares                      | Mean Square                         | F                         |
|-------------------------------------|--------------------|-------------------------------------|-------------------------------------|---------------------------|
| Fitting $\alpha$ after $\mu$        | $a - 1$            | $\text{SS}(\alpha \mid \mu)$        | $\text{MS}(\alpha \mid \mu)$        | $F(a - 1, n + 1 - a - b)$ |
| Fitting $\beta$ after $\mu, \alpha$ | $b - 1$            | $\text{SS}(\beta \mid \mu, \alpha)$ | $\text{MS}(\beta \mid \mu, \alpha)$ | $F(b - 1, n + 1 - a - b)$ |
| Error                               | $n + 1 - a - b$    | SSE                                 | MSE                                 |                           |
| Total                               | $n - 1$            | SST                                 |                                     |                           |

- $$\text{SS}(\alpha \mid \mu) = \sum_{i=1}^a \frac{Y_{i.}^2}{n_{i.}} - \frac{Y_{..}^2}{n}.$$

- $$\text{SS}(\beta \mid \mu, \alpha) = \text{SS}(\alpha, \beta \mid \mu) - \text{SS}(\alpha \mid \mu) = \vec{\beta}'_{b-1} \vec{r}.$$

### ANOVA Table III

| Source of Variation                 | Degrees of Freedom | Sum of Squares            | Mean Square               | $F$                       |
|-------------------------------------|--------------------|---------------------------|---------------------------|---------------------------|
| Fitting $\beta$ after $\mu$         | $b - 1$            | $SS(\beta   \mu)$         | $MS(\beta   \mu)$         | $F(b - 1, n + 1 - a - b)$ |
| Fitting $\alpha$ after $\mu, \beta$ | $a - 1$            | $SS(\alpha   \mu, \beta)$ | $MS(\alpha   \mu, \beta)$ | $F(a - 1, n + 1 - a - b)$ |
| Error                               | $n + 1 - a - b$    | SSE                       | MSE                       |                           |
| Total                               | $n - 1$            | SST                       |                           |                           |

- $SS(\beta | \mu) = \sum_{j=1}^b \frac{Y_{.j}^2}{n_{.j}} - \frac{Y_{..}^2}{n}$ .

- $SS(\alpha | \mu, \beta) = SS(\alpha, \beta | \mu) - SS(\beta | \mu)$ .

### Estimable Functions

$$\mathbb{E}[Y_{ij}] = \mu + \alpha_i + \beta_j = \sum_{i,j} w_{ij}(\mu + \alpha_i + \beta_j).$$

- (1)  $\mu + \alpha_i + \beta_j$  is estimable for all  $i, j$ . Choose  $w_{ij} = 1$  and  $w_{k\ell} = 0$  for  $k \neq i$  or  $\ell \neq j$ .
- (2)  $\alpha_i - \alpha_k$  is estimable for all  $i \neq k$ . Choose  $w_{ij} = -w_{kj} = 1$ , other  $w_{k\ell} = 0$ .
- (3)  $\beta_k - \beta_\ell$  is estimable for all  $k \neq \ell$ . Choose  $w_{ik} = -w_{i\ell} = 1$  and other coefficients 0.

#### EXAMPLE 18.1

Number of seconds beyond 3 minutes taken to boil 8 cups of water.

| Brand of Stove ( $\alpha$ ) | Make of Pan ( $\beta$ ) |    |    |
|-----------------------------|-------------------------|----|----|
|                             | A                       | B  | C  |
| X                           | 18                      | 12 | 24 |
| Y                           | —                       | —  | 9  |
| Z                           | 3                       | —  | 15 |
| W                           | 6                       | 3  | 18 |

The model will be  $Y_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$ .

$$\vec{Y} = (Y_{11} \ Y_{12} \ Y_{13} \ Y_{23} \ Y_{31} \ Y_{33} \ Y_{41} \ Y_{42} \ Y_{43})'$$

$$\mathbf{X} = \begin{pmatrix} \mu & \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \beta_1 & \beta_2 & \beta_3 \\ 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix} \in \mathbb{R}^{9 \times 8}$$

- $n = 9$ .
- $m = 1 + 4 + 3 = 8$ .
- $n_{..} = 9$ .
- $n_{1.} = 3, n_{2.} = 1, n_{3.} = 2, n_{4.} = 3$ .

- $n_1 = 3, n_2 = 2, n_3 = 4$ .
- $a = 4, b = 3$ , so  $a > b$  (choose  $\mu = 0, \alpha_1 = 0$ )

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} 9 & 3 & 1 & 2 & 3 & 3 & 2 & 4 \\ 3 & 3 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 2 & 0 & 0 & 2 & 0 & 1 & 0 & 1 \\ 3 & 0 & 0 & 0 & 3 & 1 & 1 & 1 \\ 3 & 1 & 0 & 1 & 1 & 3 & 0 & 0 \\ 2 & 1 & 0 & 0 & 1 & 0 & 2 & 0 \\ 4 & 1 & 1 & 1 & 1 & 0 & 0 & 4 \end{pmatrix}$$

- $\mathbf{D}_a = \text{diag}(3, 1, 2, 3)$ .

- $\mathbf{N} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{pmatrix}$ .

- $\mathbf{M} = \mathbf{D}_a^{-1}\mathbf{N} = \begin{pmatrix} 1/3 & 1/3 \\ 0 & 0 \\ 1/2 & 0 \\ 1/3 & 1/3 \end{pmatrix}$ .

- $\mathbf{D}_{(b-1)} = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}$ .

- $\mathbf{C} = \mathbf{D}_{(b-1)} - \mathbf{N}'\mathbf{D}_a^{-1}\mathbf{N} = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1/3 & 1/3 \\ 0 & 0 \\ 1/2 & 0 \\ 1/3 & 1/3 \end{pmatrix} = \begin{pmatrix} 11/6 & -4/6 \\ -4/6 & 8/6 \end{pmatrix}$ .

- $\mathbf{C}^{-1} = \frac{1}{12} \begin{pmatrix} 8 & 4 \\ 4 & 11 \end{pmatrix}$ .

- $\vec{Y}_{a.} = (18, 9, 9, 9)'$ .

- $\vec{Y}_{.a} = (54, 9, 18, 27)'$ .

- $\vec{Y}_{(b-1)} = (27, 15)'$ .

- $\vec{r} = \vec{Y}_{(b-1)} - \mathbf{M}'\vec{Y}_{a.} = \begin{pmatrix} 27 \\ 15 \end{pmatrix} - \begin{pmatrix} 1/3 & 0 & 1/2 & 1/3 \\ 1/3 & 0 & 0 & 1/3 \end{pmatrix} \begin{pmatrix} 54 \\ 9 \\ 18 \\ 27 \end{pmatrix} = \begin{pmatrix} -9 \\ -12 \end{pmatrix}$ .

Therefore, a solution is  $\vec{\beta}_0 = (0 \ 26 \ 9 \ 14 \ 17 \ -10 \ -14 \ 0)'$ . A  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \frac{1}{12} \begin{pmatrix} 0 & \begin{bmatrix} 0 & 0 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 0 \end{bmatrix} & 0 \\ \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} & \begin{bmatrix} 7 & 0 & 2 & 3 \\ 0 & 12 & 0 & 0 \\ 2 & 0 & 8 & 2 \\ 3 & 0 & 2 & 7 \end{bmatrix} & \begin{bmatrix} -4 & -5 \\ 0 & 0 \\ -4 & -2 \\ -4 & -5 \end{bmatrix} & \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \\ \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} & \begin{bmatrix} -4 & 0 & -4 & -4 \\ -5 & 0 & -2 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 8 & 4 \\ 4 & 11 \\ 0 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \end{pmatrix}.$$

## 19 Lecture 19: Two-way Crossed Classification (No Interaction) Part III

- $n_{..} = \sum_{i=1}^a \sum_{j=1}^{b_i} n_{ij}$ .
- $\vec{n}_{a.} = (n_{1.}, \dots, n_{a.})'$ .
- $\vec{n}_{.b} = (n_{.1}, \dots, n_{.b})'$ .
- $\mathbf{D}_{a.} = \text{diag}(n_{1.}, \dots, n_{a.})$ .
- $\mathbf{D}_{.b} = \text{diag}(n_{.1}, \dots, n_{.b})$ .
- $\mathbf{N} = \begin{pmatrix} n_{11} & \cdots & n_{1(b-1)} \\ \vdots & \ddots & \vdots \\ n_{a1} & \cdots & n_{a(b-1)} \end{pmatrix}$ .
- $\vec{n}_b = (n_{1b}, \dots, n_{ab})'$ .

Using these definitions, we may write

$$\mathbf{X}'\mathbf{X} = \begin{pmatrix} n_{..} & \vec{n}'_{a.} & \vec{n}'_{.(b-1)} & n_{.b} \\ \vec{n}_{a.} & \mathbf{D}_{a.} & \mathbf{N} & \vec{n}_b \\ \vec{n}_{.(b-1)} & \mathbf{N}' & \mathbf{D}_{.(b-1)} & \mathbf{O} \\ n_{.b} & \vec{n}'_b & \mathbf{O} & n_{.b} \end{pmatrix}.$$

A  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \begin{pmatrix} 0 & \mathbf{O} & \mathbf{O} & 0 \\ \mathbf{O} & \mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}' & -\mathbf{M}\mathbf{C}^{-1} & \mathbf{O} \\ \mathbf{O} & -\mathbf{C}^{-1}\mathbf{M}' & \mathbf{C}^{-1} & \mathbf{O} \\ 0 & \mathbf{O} & \mathbf{O} & 0 \end{pmatrix},$$

where the dimensions of  $\mathbf{O}$  need to be filled in for A5Q2 (trivial). Recall that  $\mathbf{M} = \mathbf{D}_{a.}^{-1}\mathbf{N}$  and  $\mathbf{C} = \mathbf{D}_{.(b-1)} - \mathbf{N}'\mathbf{D}_{a.}^{-1}\mathbf{N} = \mathbf{D}_{.(b-1)} - \mathbf{M}'\mathbf{N}$ . We will compute the following quantity.

$$\mathbf{F}\mathbf{X}'\mathbf{X} = \begin{pmatrix} \mathbf{O} & \mathbf{O} & \mathbf{O} & \mathbf{O} \\ \boxed{1} & \boxed{5} & \boxed{6} & \boxed{3} \\ \boxed{2} & \boxed{7} & \boxed{8} & \boxed{4} \\ \mathbf{O} & \mathbf{O} & \mathbf{O} & \mathbf{O} \end{pmatrix}.$$

### Block 1

$$\begin{aligned} \boxed{1} &= (\mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\vec{n}_{a.} - \mathbf{M}\mathbf{C}^{-1}\vec{n}_{.(b-1)} \\ &= \mathbf{D}_{a.}^{-1}\vec{n}_{a.} + \mathbf{M}\mathbf{C}^{-1}[\mathbf{M}'\vec{n}_{a.} - \vec{n}_{.(b-1)}] \\ &= \vec{j}_a + \mathbf{M}\mathbf{C}^{-1}\mathbf{O} \\ &= \vec{j}_a. \end{aligned}$$

$$\begin{aligned}
\mathbf{M} = \mathbf{D}_a^{-1}\mathbf{N} &\implies \mathbf{M}'\vec{n}_a = \mathbf{N}'\mathbf{D}_a^{-1}\vec{n}_a \\
&= \mathbf{N}'\vec{j}_a \\
&= \begin{pmatrix} n_{11} & \cdots & n_{1(b-1)} \\ \vdots & \ddots & \vdots \\ n_{a1} & \cdots & n_{a(b-1)} \end{pmatrix} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \\
&= \begin{pmatrix} n_{.1} \\ \vdots \\ n_{.(b-1)} \end{pmatrix}.
\end{aligned}$$

## Block 2

$$\begin{aligned}
\boxed{2} &= -\mathbf{C}^{-1}\mathbf{M}'\vec{n}_a + \mathbf{C}^{-1}\vec{n}_{.(b-1)} \\
&= -\mathbf{C}^{-1}[\mathbf{M}'\vec{n}_a - \vec{n}_{.(b-1)}] \\
&= \mathbf{O}.
\end{aligned}$$

## Block 5

$$\begin{aligned}
\boxed{5} &= (\mathbf{D}_a^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\mathbf{D}_a - \mathbf{M}\mathbf{C}^{-1}\mathbf{N}' \\
&= \mathbf{D}_a^{-1}\mathbf{D}_a + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}'\mathbf{D}_a - \mathbf{M}\mathbf{C}^{-1}\mathbf{N}' \\
&= \mathbf{I}_a + \mathbf{M}\mathbf{C}^{-1}[\mathbf{M}'\mathbf{D}_a - \mathbf{N}'] \\
&= \mathbf{I}_a.
\end{aligned}$$

$$\mathbf{M}' = \mathbf{N}'\mathbf{D}_a^{-1} \implies \mathbf{M}'\mathbf{D}_a = \mathbf{N}'\mathbf{D}_a^{-1}\mathbf{D}_a = \mathbf{N}'.$$

## Block 6

$$\begin{aligned}
\boxed{6} &= (\mathbf{D}_a^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\mathbf{N} - \mathbf{M}\mathbf{C}^{-1}\mathbf{D}_{.(b-1)} \\
&= \mathbf{D}_a^{-1}\mathbf{N} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}'\mathbf{N} - \mathbf{M}\mathbf{C}^{-1}\mathbf{D}_{.(b-1)} \\
&= \mathbf{M} + \mathbf{M}\mathbf{C}^{-1}(\mathbf{M}'\mathbf{N} - \mathbf{D}_{.(b-1)}) \\
&= \mathbf{M} - \mathbf{M}\mathbf{C}^{-1}\mathbf{C} \\
&= \mathbf{O}.
\end{aligned}$$

$$\mathbf{C} = \mathbf{D}_{.(b-1)} - \mathbf{N}'\mathbf{D}_a^{-1}\mathbf{N} = \mathbf{D}_{.(b-1)} - \mathbf{M}'\mathbf{N}.$$

## Block 7

$$\begin{aligned}
\boxed{7} &= -\mathbf{C}^{-1}\mathbf{M}'\mathbf{D}_a + \mathbf{C}^{-1}\mathbf{N}' \\
&= -\mathbf{C}^{-1}\mathbf{N}'\mathbf{D}_a^{-1}\mathbf{D}_a + \mathbf{C}^{-1}\mathbf{N}' \\
&= \mathbf{O}.
\end{aligned}$$

## Block 8

$$\begin{aligned}
\boxed{8} &= -\mathbf{C}^{-1}\mathbf{M}'\mathbf{N} + \mathbf{C}^{-1}\mathbf{D}_{.(b-1)} \\
&= -\mathbf{C}^{-1}(\mathbf{D}_{.(b-1)} - \mathbf{M}'\mathbf{N}) \\
&= \mathbf{C}^{-1}\mathbf{C} \\
&= \mathbf{I}_{b-1}.
\end{aligned}$$



### Block 3

$$\begin{aligned}
\boxed{3} &= (\mathbf{D}_{.(b-1)}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\vec{n}_b \\
&= (\mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')(\vec{n}_{a.} - \mathbf{N}\vec{j}_{b-1}) \\
&= \mathbf{D}_{a.}^{-1}\vec{n}_{a.} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}'\vec{n}_{a.} - (\mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\mathbf{N}\vec{j}_{b-1} \\
&= \vec{j}_a + \mathbf{M}\mathbf{C}^{-1}\mathbf{N}'\mathbf{D}_{a.}^{-1}\vec{n}_{a.} - (\mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\mathbf{N}\vec{j}_{b-1} \\
&= \vec{j}_a + \mathbf{M}\mathbf{C}^{-1}\vec{j}_a - \mathbf{M}\mathbf{C}^{-1}\mathbf{D}_{.(b-1)}\vec{j}_{b-1} \\
&= \vec{j}_a + \mathbf{M}\mathbf{C}^{-1}[\mathbf{N}'\vec{j}_a - \mathbf{D}_{.(b-1)}\vec{j}_{b-1}] \\
&= \vec{j}_a + \mathbf{M}\mathbf{C}^{-1}\left(\begin{pmatrix} n_{1.} \\ \vdots \\ n_{a.(b-1)} \end{pmatrix} - \begin{pmatrix} n_{1.} \\ \vdots \\ n_{.(b-1)} \end{pmatrix}\right) \\
&= \vec{j}_a.
\end{aligned}$$

$$\begin{aligned}
\vec{n}_{.a} &= \begin{pmatrix} n_{1.} \\ \vdots \\ n_{a.} \end{pmatrix} = \begin{pmatrix} n_{11} + \cdots + n_{1(b-1)} + n_{1b} \\ \vdots \\ n_{a1} + \cdots + n_{a(b-1)} + n_{ab} \end{pmatrix} \\
&= \mathbf{N}\vec{j}_{b-1} + \vec{n}_b.
\end{aligned}$$

$$\begin{aligned}
(\mathbf{D}_{a.}^{-1} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}')\mathbf{N} - \mathbf{M}\mathbf{C}^{-1}\mathbf{D}_{.(b-1)} &= \mathbf{D}_{a.}^{-1}\mathbf{N} + \mathbf{M}\mathbf{C}^{-1}\mathbf{M}'\mathbf{N} - \mathbf{M}\mathbf{C}^{-1}\mathbf{D}_{.(b-1)} \\
&= \mathbf{M} - \mathbf{M}\mathbf{C}^{-1}(\mathbf{D}_{.(b-1)} - \mathbf{M}'\mathbf{N}) \\
&= \mathbf{M} - \mathbf{M} \\
&= \mathbf{O}.
\end{aligned}$$

### Block 4

$$\begin{aligned}
\boxed{4} &= -\mathbf{C}^{-1}\mathbf{M}'\vec{n}_b \\
&= -\mathbf{C}^{-1}\mathbf{M}'[\vec{n}_{a.} - \mathbf{N}\vec{j}_{b-1}] \\
&= -\mathbf{C}^{-1}\mathbf{M}'\vec{n}_{a.} + \mathbf{C}^{-1}\mathbf{M}'\mathbf{N}\vec{j}_{b-1} \\
&= -\mathbf{C}^{-1}\mathbf{M}'\vec{n}_{a.} + (\mathbf{C}^{-1}\mathbf{D}_{.(b-1)} - \mathbf{I}_{b-1})\vec{j}_{b-1} \\
&= -\mathbf{C}^{-1}\mathbf{M}'\vec{n}_{a.} + \mathbf{C}^{-1}\mathbf{D}_{.(b-1)}\vec{j}_{b-1} - \vec{j}_{b-1} \\
&= -\mathbf{C}^{-1}[\mathbf{M}'\vec{n}_{a.} - \mathbf{D}_{.(b-1)}\vec{j}_{b-1}] - \vec{j}_{b-1} \\
&= -\mathbf{C}^{-1}\mathbf{O} - \vec{j}_{b-1} \\
&= -\vec{j}_{b-1}.
\end{aligned}$$

Therefore,

$$\mathbf{F}\mathbf{X}'\mathbf{X} = \begin{pmatrix} 0 & \mathbf{O} & \mathbf{O} & 0 \\ \vec{j}_a & \mathbf{I}_a & \mathbf{O} & \vec{j}_a \\ \mathbf{O} & \mathbf{O} & \mathbf{I}_{b-1} & -\vec{j}_{b-1} \\ 0 & \mathbf{O} & \mathbf{O} & 0 \end{pmatrix}.$$

## Estimable Combinations

$\vec{w}'\mathbf{F}\mathbf{X}'\mathbf{X} = \vec{w}$ , where  $\vec{w} = (w_0 \ w_1 \ \cdots \ w_a \ w_{a+1} \ \cdots \ w_{a+b})$ .

$$\begin{aligned}\vec{w}'\vec{\beta} &= w_0\mu + w_1\alpha_1 + \cdots + w_a\alpha_a + w_{a+1}\beta_1 + \cdots + w_{a+b}\beta_b \\ &= (w_1 + \cdots + w_a)\mu + w_1\alpha_1 + \cdots + w_a\alpha_a \\ &\quad + w_{a+1}\beta_1 + \cdots + w_{a+b-1}\beta_{b-1} \\ &\quad + [(w_1 + \cdots + w_a) - (w_{a+1} + \cdots + w_{a+b-1})]\beta_b \\ &= \sum_{i=1}^a w_i(\mu + \alpha_i + \beta_{a+b}) + \sum_{j=1}^{b-1} w_{j+a}(\beta_j - \beta_b).\end{aligned}$$

$$\vec{w}' \begin{pmatrix} 0 & \mathbf{O} & \mathbf{O} & 0 \\ \vec{j}_a & \mathbf{I}_a & \mathbf{O} & \vec{j}_a \\ \mathbf{O} & \mathbf{O} & \mathbf{I}_{b-1} & -\vec{j}_{b-1} \\ 0 & \mathbf{O} & \mathbf{O} & 0 \end{pmatrix} = \begin{pmatrix} w_1 + \cdots + w_a \\ w_1 + \cdots + w_a + w_{a+1} + \cdots + w_{a+b-1} \\ (w_1 + \cdots + w_a) - (w_{a+1} + \cdots + w_{a+b-1}) \end{pmatrix}'$$

Therefore,

$$w_0 = w_1 + \cdots + w_a.$$

$$w_{a+b} = w_1 + \cdots + w_a - (w_{a+1} + \cdots + w_{a+b-1}).$$

- (1)  $\mu + \alpha_k + \beta_\ell$  is estimable.  $w_k = w_{\ell+a} = 1$  for  $\ell < b$ , and  $w_k = 1$  otherwise.
- (2)  $\alpha_k - \alpha_\ell$  is estimable.
- (3)  $\beta_k - \beta_\ell$  is estimable.
- (4) Fix  $1 \leq k \leq b-1$ , let  $w_i = n_{ik}$ ,  $w_{a+b} = n_{.k}$ , others 0.

$$\vec{w}'\vec{\beta} = \sum n_{ik}(\mu + \alpha_i) + n_{.k}\beta_{a+b} + n_{ik}(\beta_k - \beta_{a+b}) = n_{.k}\mu + \left( n_{.k}\beta_k + \sum_{i=1}^a n_{ik}\alpha_i \right).$$

For  $k = a + b$ , choose  $w_i = n_{ik}$ ,  $w_{a+j} = 0$  for all  $j$ . Hence,

$$\vec{w}'\vec{\beta} = n_{.k}\mu + \left( n_{.k}\beta_k + \sum_{i=1}^a n_{ik}\alpha_i \right)$$

Deadly Exercise: Verify that  $\mathbf{X}'\mathbf{X}\mathbf{F}\mathbf{X}'\mathbf{X} = \mathbf{X}'\mathbf{X}$ .

LECTURE 20  
23rd March

## 20 Lecture 20: Two-way Crossed Classification (No Interaction) Part IV

$$Y_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}.$$

$\mathbf{H}_0: \mathbf{B}'\vec{\beta} = \vec{0}$  versus  $\mathbf{H}_A: \mathbf{B}'\vec{\beta} \neq \vec{0}$

The test statistic in general will be  $F = \frac{Q/r}{\text{MSE}}$ , where

- $Q = (\mathbf{B}'\vec{\beta}_0)'(\mathbf{B}'\mathbf{F}\mathbf{B})^{-1}(\mathbf{B}'\vec{\beta}_0)$ ;
- $\vec{\beta}_0$  is one solution to the normal equation  $\mathbf{X}'\mathbf{X}\vec{\beta} = \mathbf{X}'\vec{Y}$ ;
- $\mathbf{F}$  is a  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$ ;
- $r = \text{rank}(\mathbf{B})$ .

$H_0: \beta_1 = \dots = \beta_b$  versus  $H_A: \neg H_0$

$$\bullet \mathbf{B}' = \begin{pmatrix} \mu & \alpha_1 & \dots & \alpha_a & \beta_1 & \beta_2 & \dots & \beta_{b-1} & \beta_b \\ 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 & -1 \\ 0 & 0 & \dots & 0 & 0 & 1 & \ddots & 0 & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 & \dots & 0 & 1 & -1 \end{pmatrix} = (\mathbf{O}_{(b-1) \times (a+1)} \quad \mathbf{I}_{b-1} \quad -\vec{j}_{b-1}).$$

- A5Q2: Show that  $Q = SS(\beta \mid \mu, \alpha)$ , and  $r = b - 1$ .

$H_0: \alpha_1 = \dots = \alpha_a$  versus  $H_A: \neg H_0$

- $\mathbf{B}' = (0 \quad \mathbf{I}_{a-1} \quad -\vec{j}_{a-1} \quad \mathbf{O}_{(a-1) \times b})$ .
- $Q = SS(\alpha \mid \mu, \beta)$  and  $r = a - 1$ .

$H_0: \beta_j + \frac{1}{n_j} \sum_{i=1}^a n_{ij} \alpha_i$  are equal for  $j = 1, \dots, b$  versus  $H_A: \neg H_0$

Idea: The first equation will be

$$\beta_1 + \frac{1}{n_{.1}} \sum_{i=1}^a n_{i1} \alpha_i = \beta_b + \frac{1}{n_{.b}} \sum_{i=1}^a n_{ib} \alpha_i \implies \beta_1 - \beta_b + \sum_{i=1}^a \left( \frac{n_{i1}}{n_{.1}} - \frac{n_{ib}}{n_{.b}} \right) \alpha_i = 0.$$

$$\bullet \mathbf{B}' = \begin{pmatrix} \mu & \alpha_1 & \dots & \alpha_a & \beta_1 & \beta_2 & \dots & \beta_{b-1} & \beta_b \\ 0 & \frac{n_{11}}{n_{.1}} - \frac{n_{1b}}{n_{.b}} & \dots & \frac{n_{a1}}{n_{.1}} - \frac{n_{ab}}{n_{.b}} & 1 & 0 & \dots & 0 & -1 \\ 0 & \frac{n_{12}}{n_{.2}} - \frac{n_{1b}}{n_{.b}} & \dots & \frac{n_{a2}}{n_{.2}} - \frac{n_{ab}}{n_{.b}} & 0 & 1 & \ddots & 0 & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \end{pmatrix}.$$

- $Q = SS(\beta \mid \mu)$  and  $r = b - 1$ .

Notes:

$$\frac{n_{11}}{n_{.1}} + \dots + \frac{n_{a1}}{n_{.1}} = 1.$$

$$\beta_1 + \frac{n_{11}}{n_{.1}} \alpha_1 + \dots + \frac{n_{a1}}{n_{.1}} \alpha_a = \beta_b + \frac{n_{1b}}{n_{.b}} \alpha_1 + \dots + \frac{n_{ab}}{n_{.b}} \alpha_a.$$

If  $n_{ij} = 1$  for all  $i = 1, \dots, a$  and  $j = 1, \dots, b$ , then we will be testing  $\beta_1 = \beta_b$ .

$H_0: \alpha_i + \frac{1}{n_i} \sum_{j=1}^b n_{ij} \beta_j$  are equal for  $i = 1, \dots, a$  versus  $H_A: \neg H_0$

- $\mathbf{B}' =$  exercise.
- $Q = SS(\alpha \mid \mu)$  and  $r = a - 1$ .

## Existence of Interaction

### EXAMPLE 20.1

No interaction:

| Age \ Type | A   | B   | C   |
|------------|-----|-----|-----|
| <30        | 3.2 | 1.8 | 6.5 |
| 30-60      | 2.8 | 1.6 | 5.3 |
| >60        | 1.5 | 1.0 | 3.4 |

Interaction:

| Age \ Type | A   | B   | C   |
|------------|-----|-----|-----|
| <30        | 3.2 | 1.4 | 6.5 |
| 30-60      | 6.2 | 1.2 | 2.5 |
| >60        | 5.8 | 1.3 | 2.1 |

Look at the plots.

## Two-Way Crossed Classification (Interaction) Part I

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk},$$

where  $i = 1, \dots, a; j = 1, \dots, b; k = 1, \dots, n_{ij}$  where  $n_{ij} \geq 1$ .

| $i \setminus j$ | 1        | 2        | ...      | $b$      |
|-----------------|----------|----------|----------|----------|
| 1               | $n_{11}$ | $n_{12}$ | ...      | $n_{1b}$ |
| 2               | $n_{21}$ | $n_{22}$ | ...      | $n_{2b}$ |
| $\vdots$        | $\vdots$ | $\vdots$ | $\ddots$ | $\vdots$ |
| $a$             | $n_{a1}$ | $n_{a2}$ | ...      | $n_{ab}$ |

$$Y_{i.j.} = \sum_{k=1}^{n_{ij}} Y_{ijk} \implies \bar{Y}_{ij.} = \frac{1}{n_{ij}} Y_{i.j.}$$

$$Y_{i..} = \sum_{j=1}^b \sum_{k=1}^{n_{ij}} Y_{ijk},$$

$$Y_{.j.} = \sum_{i=1}^a \sum_{k=1}^{n_{ij}} Y_{ijk},$$

$$n_{i.} = \sum_{j=1}^b n_{ij},$$

$$n_{.j} = \sum_{i=1}^a n_{ij},$$

$$n_{..} = \sum_{i=1}^a \sum_{j=1}^b n_{ij},$$

$s =$  total number of non-empty cells.

The design matrix  $\mathbf{X}$  is  $n_{..} \times m$ , where  $m = 1 + a + b + s$  and  $\text{rank}(\mathbf{X}'\mathbf{X}) = s$ .

$$\mathbf{D} = \text{diag}\left(\frac{1}{n_{11}}, \dots, \frac{1}{n_{1b}}, \frac{1}{n_{21}}, \dots, \frac{1}{n_{a1}}, \dots, \frac{1}{n_{ab}}\right) \in \mathbb{R}^{s \times s},$$

where  $n_{ij} \geq 1$ . A  $g$ -inverse of  $\mathbf{X}'\mathbf{X}$  is

$$\mathbf{F} = \begin{pmatrix} \mathbf{O} & \mathbf{O} \\ \mathbf{O} & \mathbf{D} \end{pmatrix} \in \mathbb{R}^{m \times m}.$$

A solution to the normal equation is

$$\vec{\beta}_0 = \mathbf{F}\mathbf{X}'\vec{Y} = (\mathbf{O}_{1 \times (1+a+b)} \quad \bar{Y}_{11.} \quad \dots \quad \bar{Y}_{ab.}),$$

where we only keep the non-empty cells (i.e.,  $n_{ij} \geq 1$ ). We can test  $H_0: \mathbf{B}'\vec{\beta} = \vec{m}$  versus  $\mathbf{B}'\vec{\beta} \neq \vec{m}$ .

- $SST = \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^{n_{ij}} Y_{ijk}^2 - \frac{Y_{\dots}^2}{n_{\dots}}$ .
- $SSE = \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^{n_{ij}} Y_{ijk}^2 - \sum_{i=1}^a \sum_{j=1}^b \frac{Y_{ij.}^2}{n_{ij}}$ .

### ANOVA Table I

| SV                               | df      | SS                                   | MS                                   | F                                   |
|----------------------------------|---------|--------------------------------------|--------------------------------------|-------------------------------------|
| $\alpha, \beta, \gamma \mid \mu$ | $s - 1$ | $SS(\alpha, \beta, \gamma \mid \mu)$ | $MS(\alpha, \beta, \gamma \mid \mu)$ | $F(\alpha, \beta, \gamma \mid \mu)$ |
| Error                            | $n - s$ | SSE                                  | MSE                                  |                                     |
| Total                            | $n - 1$ | SST                                  |                                      |                                     |

- $SS(\alpha, \beta, \gamma \mid \mu) = \sum_{i=1}^a \sum_{j=1}^b \frac{Y_{ij.}^2}{n_{ij}} - \frac{Y_{\dots}^2}{n_{\dots}}$

### ANOVA Table II

Fitting  $\alpha$ , then  $\beta$ , followed by  $\gamma$ . In compact notation,  $\alpha \rightarrow \beta \rightarrow \gamma$ .

| SV                               | df              | SS                                   | MS                                   | F                                   |
|----------------------------------|-----------------|--------------------------------------|--------------------------------------|-------------------------------------|
| $\alpha \mid \mu$                | $a - 1$         | $SS(\alpha \mid \mu)$                | $MS(\alpha \mid \mu)$                | $F(\alpha \mid \mu)$                |
| $\beta \mid \mu, \alpha$         | $b - 1$         | $SS(\beta \mid \mu, \alpha)$         | $MS(\beta \mid \mu, \alpha)$         | $F(\beta \mid \mu, \alpha)$         |
| $\gamma \mid \mu, \alpha, \beta$ | $s - a - b + 1$ | $SS(\gamma \mid \mu, \alpha, \beta)$ | $MS(\gamma \mid \mu, \alpha, \beta)$ | $F(\gamma \mid \mu, \alpha, \beta)$ |
| Error                            | $n - s$         | SSE                                  | MSE                                  |                                     |
| Total                            | $n - 1$         | SST                                  |                                      |                                     |

- $SS(\alpha \mid \mu) = \sum_{i=1}^a \frac{Y_{i..}^2}{n_i} - \frac{Y_{\dots}^2}{n_{\dots}}$ .
- $SS(\beta \mid \mu, \alpha) = (\vec{\beta}_{b-1}^0)' \vec{r}$ .
- $SS(\gamma \mid \mu, \alpha, \beta) = \sum_{i=1}^a \sum_{j=1}^b \frac{Y_{ij.}^2}{n_{ij}} - \sum_{i=1}^a \frac{Y_{i..}^2}{n_i} - (\vec{\beta}_{b-1}^0)' \vec{r}$ , where

$$\vec{\beta}_0 = (\mu^0 \quad \alpha_1^0 \quad \dots \quad \alpha_a^0 \quad \beta_1^0 \quad \dots \quad \beta_b^0 \quad (\gamma_{ij})^0)'$$

$$\vec{\beta}_{b-1}^0 = (\beta_1^0 \quad \dots \quad \beta_{b-1}^0)'$$

To obtain  $\vec{\beta}_{b-1}^0$  and  $\vec{r}$ , we replace  $Y_{.j}$  and  $\bar{Y}_{i.}$  by  $Y_{.j}$  and  $\bar{Y}_{i.}$ , respectively.

## 21 Lecture 21: Two-way Crossed Classification (Interaction) Part II

### Estimable Combinations

$\vec{b}'\vec{\beta}$  is estimable if and only if  $\vec{b}' = \vec{w}'\mathbf{X}$ . Therefore,  $\vec{w}'\mathbf{X}\vec{\beta}$  is always estimable.

$$\vec{\beta}' = (\mu, \alpha_1, \dots, \alpha_a, \beta_1, \dots, \beta_b, \{\gamma_{ij}\})'$$

where we only include the non-empty  $(i, j)$  in  $\{\gamma_{ij}\}$ .

$$(\mathbf{X}\vec{\beta})' = \underbrace{(\mu + \alpha_1 + \beta_1 + \gamma_{11}, \dots, \mu + \alpha_1 + \beta_1 + \gamma_{11}, \dots)}_{n_{11}}, \dots, \underbrace{(\mu + \alpha_a + \beta_b + \gamma_{ab}, \dots, \mu + \alpha_a + \beta_b + \gamma_{ab})}_{n_{ab}}'.$$

$$\vec{w}'\mathbf{X}\vec{\beta} = \sum_{i=1}^a \sum_{j=1}^b c_{ij} \mu_{ij},$$

where  $\mu_{ij} = \mu + \alpha_i + \beta_j + \gamma_{ij}$ .

- (1)  $\mu_{ij} = \mu + \alpha_i + \beta_j + \gamma_{ij}$  is estimable.
- (2) For any  $k, \ell$ , define

$$\Lambda_{k\ell} = \alpha_k + \frac{1}{n_k} \sum_{j=1}^b n_{kj}(\beta_j + \gamma_{kj}) - \left[ \alpha_\ell + \frac{1}{n_\ell} \sum_{j=1}^b n_{\ell j}(\beta_j + \gamma_{\ell j}) \right].$$

Claim:  $\Lambda_{k\ell}$  is estimable.

First term,

$$\begin{aligned} \alpha_k + \frac{1}{n_k} \sum_{j=1}^b n_{kj}(\beta_j + \gamma_{kj}) &= \frac{1}{n_k} \sum_{j=1}^b n_{kj} \alpha_k + \frac{1}{n_k} \sum_{j=1}^b n_{kj}(\beta_j + \gamma_{kj}) \\ &= \frac{1}{n_k} \sum_{j=1}^b n_{kj}(\alpha_k + \beta_k + \gamma_{kj} + \mu - \mu) \\ &= -\mu + \frac{1}{n_k} \sum_{j=1}^b n_{kj}(\mu + \alpha_k + \beta_j + \gamma_{kj}). \end{aligned}$$

Similarly,

$$\alpha_\ell + \frac{1}{n_\ell} \sum_{j=1}^b n_{\ell j}(\beta_j + \gamma_{\ell j}) = -\mu + \frac{1}{n_\ell} \sum_{j=1}^b n_{\ell j}(\mu + \alpha_\ell + \beta_j + \gamma_{\ell j}).$$

Taking the difference of the first term and second term yields the result.

- (3)

$$\psi_j = \left( n_j - \sum_{i=1}^a \frac{n_{ij}^2}{n_i} \right) \beta_j - \sum_{\ell \neq j} \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} \beta_\ell + \sum_{i=1}^a \left( n_{ij} - \frac{n_{ij}^2}{n_i} \right) \gamma_{ij} - \sum_{\ell \neq j} \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} \gamma_{i\ell}.$$

Claim:  $\psi_j$  is estimable.

$$\begin{aligned} \psi_j &= \sum_{i=1}^a \left( n_{ij} - \frac{n_{ij}^2}{n_i} \right) \beta_j - \sum_{\ell \neq j} \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} \beta_\ell + \sum_{i=1}^a \left( n_{ij} - \frac{n_{ij}^2}{n_i} \right) \gamma_{ij} - \sum_{\ell \neq j} \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} \gamma_{i\ell} \\ &= \sum_{i=1}^a \left( n_{ij} - \frac{n_{ij}^2}{n_i} \right) (\beta_j + \gamma_{ij}) - \sum_{\ell \neq j} \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} (\beta_\ell + \gamma_{i\ell}) \\ &= \sum_{i=1}^a n_{ij} (\beta_j + \gamma_{ij}) - \sum_{\ell=1}^b \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} (\beta_\ell + \gamma_{i\ell}) \\ &= \sum_{i=1}^a n_{ij} (\mu + \alpha_i + \beta_j + \gamma_{ij}) - \sum_{\ell=1}^b \sum_{i=1}^a \frac{n_{ij} n_{i\ell}}{n_i} (\mu + \alpha_i + \beta_\ell + \gamma_{i\ell}), \end{aligned}$$

since

$$\begin{aligned}\sum_{i=1}^a n_{ij}(\mu + \alpha_i) &= n_{.j}\mu + \sum_{i=1}^a n_{.j}\alpha_i, \\ \sum_{\ell=1}^b \sum_{i=1}^a \frac{n_{ij}n_{i\ell}}{n_i}(\mu + \alpha_i) &= \sum_{i=1}^a (\mu + \alpha_i) \sum_{\ell=1}^b \frac{n_{ij}n_{i\ell}}{n_i} \\ &= \sum_{i=1}^a (\mu + \alpha_i)n_{ij}.\end{aligned}$$

(4) For any  $(i, j)$  and  $(k, \ell)$  let

$$\Theta_{(i,j),(k,\ell)} = \gamma_{ij} - \gamma_{i\ell} - \gamma_{kj} + \gamma_{k\ell}.$$

Claim:  $\Theta_{(i,j),(k,\ell)}$  is estimable. Proof by picture: a square. By definition,

$$\mu_{ij} = \mu + \alpha_i + \beta_j + \gamma_{ij}.$$

$$\mu_{i\ell} = \mu + \alpha_i + \beta_\ell + \gamma_{i\ell}.$$

$$\mu_{kj} = \mu + \alpha_k + \beta_j + \gamma_{kj}.$$

$$\mu_{k\ell} = \mu + \alpha_k + \beta_\ell + \gamma_{k\ell}.$$

Hence,

$$\mu_{ij} - \mu_{i\ell} = \beta_j - \beta_\ell + \gamma_{ij} - \gamma_{i\ell}.$$

$$\mu_{kj} - \mu_{k\ell} = \beta_j - \beta_\ell + \gamma_{kj} - \gamma_{k\ell}.$$

Taking the difference of the first term and second term yields the result.

## Hypothesis Testing

(1)  $H_0: \frac{1}{n_i} \sum_{j=1}^b n_{ij}\mu_{ij}$  are the same for all  $i$ .

$$\frac{n_{i1}}{n_i}\mu_{i1} + \cdots + \frac{n_{ib}}{n_i}\mu_{ib}.$$

Intuitively, we are testing the weighted means are all the same in each row.  $Q = \text{SS}(\alpha \mid \mu)$ .

$H_0: \frac{1}{n_{.j}} \sum_{i=1}^a n_{ij}\mu_{ij}$  are the same for all  $j$ .  $Q = \text{SS}(\beta \mid \mu)$ .

(2)  $H_0: \psi_j = 0$  for all  $j \leq b-1$ .  $Q = \text{SS}(\beta \mid \mu, \alpha)$ .

$$\varphi_i = \left( n_i - \sum_{j=1}^b \frac{n_{ij}^2}{n_{.j}} \right) \alpha_i - \sum_{k \neq \ell} \left( \sum_{j=1}^b \frac{n_{ij}n_{kj}}{n_{.j}} \right) \alpha_k + \sum_{j=1}^b \left( n_{ij} - \frac{n_{ij}^2}{n_{.j}} \right) \gamma_{ij} - \sum_{k \neq i} \left( \sum_{j=1}^b \frac{n_{ij}n_{kj}}{n_{.j}} \right) \gamma_{kj}.$$

$H_0: \varphi_i = 0$  for all  $i = 1, \dots, a-1$ .  $Q = \text{SS}(\alpha \mid \mu, \beta)$ . The reason we are not testing all  $i, j$  since

$$\sum_{j=1}^b \psi_j = \sum_{i=1}^a \varphi_i = 0.$$

We will show  $\sum_{i=1}^a \varphi_i = 0$ , the other is an exercise.

$$\begin{aligned} \sum_{i=1}^a \varphi_i &= \sum_{i=1}^a n_i \alpha_i - \sum_{i=1}^a \sum_{k=1}^a \sum_{j=1}^b \frac{n_{ij} n_{kj}}{n_{.j}} \alpha_k + \sum_{i=1}^a \sum_{j=1}^b n_{ij} \gamma_{ij} - \sum_{i=1}^a \sum_{k=1}^a \sum_{j=1}^b \frac{n_{ij} n_{kj}}{n_{.j}} \gamma_{kj} \\ &= \sum_{i=1}^a n_i \alpha_i - \sum_{k=1}^a \alpha_k \sum_{j=1}^b \sum_{i=1}^a \frac{n_{ij} n_{kj}}{n_{.j}} + \sum_{i=1}^a \sum_{j=1}^b n_{ij} \gamma_{ij} - \sum_{k=1}^a \sum_{j=1}^b \gamma_{kj} \sum_{i=1}^a \frac{n_{ij} n_{kj}}{n_{.j}} \\ &= \sum_{i=1}^a n_i \alpha_i - \sum_{k=1}^a n_k \alpha_k + \sum_{i=1}^a \sum_{j=1}^b n_{ij} \gamma_{ij} - \sum_{k=1}^a \sum_{j=1}^b n_{kj} \gamma_{kj} \\ &= 0. \end{aligned}$$

(3) Let  $g_1, \dots, g_{s-a-b+1}$  be linearly independent functions of  $\{\Theta_{(i,j),(k,\ell)} : (i,j), (k,\ell)\}$ .

$H_0: g_\ell = 0$  for all  $\ell = 1, \dots, s-a-b+1$ .  $Q = SS(\gamma \mid \mu, \alpha, \beta)$ .

LECTURE 22  
30th March

## 22 Lecture 22: Factorial Design

- Level of Factor I:  $a$ ;
- Level of Factor II:  $b$ ;
- Replications:  $n$  (i.e., balanced);
- Total number of observations:  $nab$ .

### EXAMPLE 22.1: $2^2$

Consider the effect on the conversion in a chemical process of the *concentration of the reactor* and the *amount of catalyst*. Each factor has two levels: low (–) and high (+).

| catalyst \ concentration | –  | +   |
|--------------------------|----|-----|
| –                        | -- | --+ |
| +                        | +- | ++  |

Replication:

|    | 1  | 2  | 3  | Total |
|----|----|----|----|-------|
| -- | 28 | 25 | 27 | 80    |
| -+ | 36 | 32 | 32 | 100   |
| +- | 18 | 19 | 23 | 60    |
| ++ | 31 | 30 | 29 | 90    |

Let  $(1) = --- = 80$ ,  $a = ++ = 60$ ,  $b = +- = 100$ , and  $ab = ++ = 90$ .

$$A = \frac{ab - b + a - (1)}{2n} \rightarrow \text{main effect of the catalyst.}$$

$$B = \frac{ab - a + b - (1)}{2n} \rightarrow \text{main effect of the concentration.}$$

$$AB = \frac{ab - b - (a - (1))}{2n} = \frac{ab - a - (b - (1))}{2n} \rightarrow \text{interaction.}$$

$$\text{Contrasts}_A = (ab + a) - (b + (1)).$$

$$\text{Contrasts}_B = (ab + b) - ((1) + a).$$

$$\text{Contrasts}_{AB} = (ab - a) - (b - (1)).$$



$Y_{ijk}$ , where  $i = 1, \dots, a$ ;  $j = 1, \dots, b$ ;  $k = 1, \dots, n$ . We can define  $Y_{\dots}$ ,  $Y_{i..}$ , and  $Y_{.j}$  as before.

$$Y_{ijk} - \bar{Y}_{\dots} = (\bar{Y}_{i..} - \bar{Y}_{\dots}) + (\bar{Y}_{.j} - \bar{Y}_{\dots}) + (\bar{Y}_{ij.} - \bar{Y}_{i..} - \bar{Y}_{.j} + \bar{Y}_{\dots}) + (Y_{ijk} - \bar{Y}_{ij.}).$$

Squaring and summing yields SST, where we note that the cross-terms will be zero. Hence,

$$\bullet \text{ SST} = \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^n Y_{ijk}^2 - \frac{Y_{\dots}^2}{nab}.$$

$$\bullet \text{ SS}_A = \frac{1}{bn} \sum_{i=1}^a Y_{i..}^2 - \frac{Y_{\dots}^2}{nab}.$$

$$\bullet \text{ SS}_B = \frac{1}{an} \sum_{j=1}^b Y_{.j}^2 - \frac{Y_{\dots}^2}{nab}.$$

$$\bullet \text{ SS}_{AB} = \frac{1}{n} \sum_{i=1}^a \sum_{j=1}^b Y_{ij.}^2 - \frac{Y_{\dots}^2}{nab}.$$

| Source of Variation | Degrees of Freedom | Sum of Squares   | Mean Square      |
|---------------------|--------------------|------------------|------------------|
| Factor A            | $a - 1$            | $\text{SS}_A$    | $\text{MS}_A$    |
| Factor B            | $b - 1$            | $\text{SS}_B$    | $\text{MS}_B$    |
| Factor AB           | $(a - 1)(b - 1)$   | $\text{SS}_{AB}$ | $\text{MS}_{AB}$ |
| Error               | $ab(n - 1)$        | $\text{SSE}$     | $\text{MSE}$     |
| Total               | $n - 1$            | $\text{SST}$     |                  |

In our case,  $a = b = 2$ .

$$\text{SS}_A = \frac{1}{4n} \left[ 2 \sum_{i=1}^n Y_{i..}^2 - Y_{\dots}^2 \right] = \frac{1}{4n} (Y_{1..} - Y_{2..})^2.$$

We can also write  $Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_1 x_2 + \varepsilon$ , where  $x_1 = \pm 1$ ,  $x_2 = \pm 1$ , and  $x_1 x_2 = \pm 1$ .

## Generalized Linear Models

In our usual case,  $\vec{Y} = \mathbf{X}\vec{\beta}$  and we assume  $\mathbb{E}[\vec{Y}] = \mathbf{X}\vec{\beta}$ . However, suppose we have  $\vec{\mu} = \mathbb{E}[\vec{Y}] = H(\mathbf{X}\vec{\beta})$ , where  $g = H^{-1}$  and  $g(\vec{\mu}) = \mathbf{X}\vec{\beta}$ .  $g$  must be monotone and differentiable.

### Exponential Family

Let  $Y$  be a random variable with pmf or pdf  $f(y, \theta)$ .  $Y$  is in the exponential family if

$$f(y, \theta) = \exp\{a(y)b(\theta) + c(\theta) + d(y)\}.$$

The distribution of  $Y$  is in canonical form if  $a(y) = y$ . Define  $\ell(y, \theta) = \log(f(y, \theta))$ .  $\frac{d\ell}{d\theta} = \frac{f'(y, \theta)}{f(y, \theta)} = U$  is called the **score function**.  $\text{Var}(U)$  is called Fisher information.

#### REMARK — (Proposition 22.1)

Assume that  $b(\cdot)$  and  $c(\cdot)$  are second order differentiable. We have

$$\mathbb{E}[a(Y)] = -\frac{c'(\theta)}{b'(\theta)}, \quad \text{Var}(a(Y)) = \frac{b''(\theta)c'(\theta) - c''(\theta)b'(\theta)}{[b'(\theta)]^3}.$$

**Proof:** Assume that  $Y$  is continuous. We know that

$$\int f(y, \theta) dy = 1 \implies \int \frac{d}{dy} f(y, \theta) dy = 0.$$

So,

$$\int f(y, \theta)[a(y)b'(\theta) + c'(\theta)] dy = 0 \implies c'(\theta) + b'(\theta) \underbrace{\int a(y)f(y, \theta) dy}_{\mathbb{E}[a(Y)]} = 0.$$

Rearranging yields the expectation. For the variance

$$\begin{aligned} 0 &= \int \frac{d^2}{d\theta^2} f(y, \theta) dy \\ &= \int \frac{d}{d\theta} \left( f(y, \theta)[a(y)b'(\theta) + c'(\theta)] \right) dy \\ &= \int f(y, \theta)[a(y)b'(\theta) + c'(\theta)]^2 dy + \int f(y, \theta)[a(y)b''(\theta) + c''(\theta)] dy \\ &= [b'(\theta)]^2 \mathbb{E}[a^2(Y)] + (2b'(\theta)c'(\theta) + b''(\theta)) \mathbb{E}[a(Y)] + [c'(\theta)]^2 + c''(\theta). \end{aligned}$$

Use the formula  $\mathbb{E}[a^2(Y)] = \text{Var}(a(Y)) + \mathbb{E}[a(Y)]^2$ , rearrange and do a bunch of algebra (the usual) to get the variance.

#### REMARK — (Proposition 22.2)

- $\mathbb{E}[U] = 0$ ;
- $\text{Var}(U) = \mathbb{E} \left[ \left( \frac{d\ell}{d\theta} \right)^2 \right] = -\mathbb{E} \left[ \frac{d^2\ell}{d\theta^2} \right] = -\mathbb{E}[U']$ .

**Proof:**

$$U = \frac{d\ell}{d\theta} = \frac{f'}{f} \implies \mathbb{E}[U] = \int \frac{f'}{f} f dy = \int f' dy = 0.$$

$$\begin{aligned} \text{Var}(U) &= \mathbb{E}[U^2] \\ &= \mathbb{E} \left[ \left( \frac{d\ell}{d\theta} \right)^2 \right] \\ &= \int \left( \frac{f'}{f} \right)^2 f dy \\ &= \int \frac{(f')^2}{f} dy - \underbrace{\int f'' dy}_0 \\ &= \int \frac{(f')^2 - f''f}{f} dy \\ &= -\int \left( \frac{f'}{f} \right)' dy \\ &= -\int \frac{d^2\ell}{d\theta^2} dy. \end{aligned}$$

## REMARK

For exponential family,

$$\begin{aligned}\text{Var}(U) &= -\mathbb{E}[a(Y)b''(\theta) + c''(\theta)] \\ &= -\mathbb{E}[a(Y)]b''(\theta) - c''(\theta) \\ &= \frac{c'}{b'}b'' - c'' \\ &= \frac{b''(\theta)c'(\theta) - c''(\theta)b'(\theta)}{b'(\theta)}.\end{aligned}$$

Therefore,  $\text{Var}(U) = [b'(\theta)]^2 \text{Var}(a(X))$ .

## DEFINITION 22.1: Generalized Linear Model

Let  $Y_1, \dots, Y_N$  be independent with the same type of distribution.

- (1) The distribution of each  $Y_i$  is in the exponential family with canonical form.
- (2)  $g$  is a monotone differentiable function such that

$$g(\mu_i) = \sum_{k=1}^r X_{ik}\beta_k,$$

and  $\mu_i = \mathbb{E}[Y_i]$  for  $i = 1, \dots, r$ , where  $g$  is called a **link function**. We must have  $r \ll n$ .

Goal: Estimation of  $r, \beta_1, \dots, \beta_r$ . We define the **explanatory matrix** as

$$\begin{pmatrix} X_{11} & \cdots & X_{1r} \\ X_{21} & \cdots & X_{2r} \\ \vdots & \ddots & \vdots \\ X_{n1} & \cdots & X_{nr} \end{pmatrix}.$$

## EXAMPLE 22.2

Suppose  $Y \sim \text{POI}(\lambda)$ .  $Y = \mathbb{E}[Y] + \text{noise}$ , hence  $\mathbb{E}[Y] = \lambda$ , where  $\text{noise} \in \mathbb{N}$ , but this does not make sense. The reason is because we cannot write  $\lambda = \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_r x_r$  since  $\lambda > 0$ . Hence, we instead consider

$$\log(\lambda) = \beta_1 x_1 + \cdots + \beta_r x_r \implies \lambda = \exp\{\beta_1 x_1 + \cdots + \beta_r x_r\}.$$

EXTRA  
3rd to 10th April

These three days were used for review and office hours.